

Note on the occurrence of phytoplankton and its relation with mass mortality in the Jakarta Bay, may and november 2004

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=118308&lokasi=lokal>

Abstrak

There were noted two times (May and November 2004), fish mass mortality occurred in the coast of the Jakarta Bay, killing several species of small and big fishes, crustaceans, eels and molluscs. Phytoplankton blooms in the Jakarta Bay is accelerated by the high input of nutrients, which caused eutropication of the Bay water. Analysis of water sample showed in st 1 (Marina coast) showed that phytoplankton abundance was of 2.9×10^6 cells/l-1, the common diatom species recorded at that time were: *Skeletonema costatum*, *Thalassiosira mala*, *Bacteriastrum varians*, *Chaetoceros pseudocurvisetus*, *Nitzschia sigma*, *Coscinodiscus radiatus*. Of the which: *Skeletonema costatum* 1.8×10^6 cells/l-1; in st. 3 Binaria lake was the most a common diatoms species found in the Jakarta Bay. This species frequently bloom, especially after rainfall, causing no harm to marine life, the second diatoms *Thalassiosira mala* 2.8×10^6 cells/l-1 (st 4 Carnaval coast), during this decade in more often frequencies. The second largest group was represented by four dinoflagellates species: *Prorocentrum micans*, *Protoperidinium* sp., *Ceratium furca* and *Gonyaulax* sp., and one species dominated of dinoflagellate *Prorocentrum micans* 2.3×10^6 cells/l-1 (in the Dadap coast). *P. micans* is also common species of the Jakarta Bay. Hydrological conditions in May and November 2004 are, temperature 32 – 33 oC, pH 7.76 to 7.92, Salinity 32 ‰, Dissolved oxygen 3.88 to 4.26 mg/l, Phosphate 0.10 – 0.40 µg-at-P l-1, Nitrate 0.01 to 0.03 µg-at-N l-1. The result suggests that phytoplankton distributes in wider areas than expected and monitoring of its occurrence in Indonesian waters is necessary to prevent harmful effects of such bloom.