

Pengembangan metode elisa langsung untuk penentuan residu kloramfenikol

Abdul Azis, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20176225&lokasi=lokal>

Abstrak

ABSTRAK

The existence of chloramphenicol residues in veterinary products could cause such a serious side effect like aplastic anemia. A specific and sensitive ELISA method to detect chloramphenicol residues was needed to be developed in order to get a more efficient and economical analysing method. In this study, a direct ELISA method has been done using chloramphenicol-bovine serum albumin (CAP-BSA) as antigen and antisera chloramphenicol labelled by enzyme horseradish peroxidase (HRP) as antibody. CAP-BSA with a dilution of 1:200 and antisera chloramphenicol labelled by HRP with a dilution of 1:125 showed a limit detection of 0.05 ng/ml. CAP-BSA had been synthesized from chloramphenicol sodium succinate and BSA, using mixed anhydride reaction, whereas HRP-labelled antisera chloramphenicol had been synthesized from antisera chloramphenicol and HRP, using periodate conjugation technique. The production of antisera chloramphenicol has been performed through CAP-BSA immunization using two New Zealand White rabbits resulting in antisera chloramphenicol with a less satisfactory specificity and sensitivity. The recovery assay of chloramphenicol in cow milk gave 98,17% \pm 2,88% yield.

<hr>