

## Uji Efek Hipoglikemik Obat Herbal FAD terhadap Tikus Putih Jantan yang Dibebani Glukosa

Agung Setiayanti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20181198&lokasi=lokal>

---

### Abstrak

Obat herbal "FAD" sebagai salah satu produk obat tradisional yang memiliki indikasi sebagai antidiabetes perlu diuji khasiatnya secara ilmiah untuk membuktikan efektifitasnya. Penelitian ini bertujuan untuk mengetahui kemampuan obat herbal "FAD" dalam menurunkan kadar glukosa darah tikus putih jantan yang dibebani glukosa. Penelitian ini dilakukan dengan rancangan acak lengkap menggunakan 30 ekor tikus putih jantan galur Sprague-Dawley yang terbagi dalam 5 kelompok. Larutan uji diberikan secara oral dengan variasi dosis, yaitu dosis 1 (216,1 mg/200 g bb tikus), dosis 2 (432,2 mg / 200g bb tikus), dan dosis 3 (864,4 mg/200 g bb tikus), sebagai kontrol pembanding digunakan suspensi metformin hidroklorida dalam CMC 0,5% 270 mg/200 g bb tikus sedangkan kontrol normal dan kontrol perlakuan diberikan suspensi CMC 0,5%. Satu jam setelah perlakuan, masing-masing tikus diberikan glukosa kecuali kontrol normal sebesar 375 mg/200g bb tikus per oral. Darah diambil pada menit ke 0, 60, 90, 120, 150 dan 180 dihitung saat setelah perlakuan. Kadar glukosa darah ditetapkan dengan metode otoluidin. Hasil penelitian menunjukkan bahwa obat herbal "FAD" dosis 864,4 mg/200 g bb tikus dapat menurunkan kadar glukosa darah pada tikus putih jantan yang dibebani glukosa.

<hr>Herbal medicine "FAD" is one of traditional medicine which has indication as antidiabetic agent need to find its effectiveness. This research was carried out to investigate reduction of blood glucose level by herbal medicine "FAD" on glucose-induced male rats. This study was conducted by employing a complete random design, using 30 male Sprague-Dawley rats which divided into 5 groups. The test solution was given by orally with variant of dose of 216,1, 432,2, 864,4 mg/200 g body weight of rat, as compared control is used metformin hidroklorida suspension in 0,5% CMC with dose of 270 mg/200 g body weight of rat, while normal control and treatment control were given 0,5% CMC suspension. One hour later, each rat was given glucose with dose 375 mg/200 g body weight of rat orally. Blood was collected at 0, 60, 90, 120, 150 and 180 minutes. Glucose blood level was determined by otoluidin method. The result of the research shows that herbal medicine "FAD" with dose 864,4 mg/200 g body weight of rat can reduce blood glucose level on glucose-induced male rats.