

Uji efektifitas gastroprotektif kombinasi ekstrak rimpang kunyit (*Curcuma domestica* Val.) dan kulit batang mimba (*Azadirachta indica* A. Juss) pada tikus putih yang diinduksi dengan asetosal

Dini Kusumaningtyas, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20181404&lokasi=lokal>

Abstrak

Beberapa penelitian melaporkan bahwa ekstrak rimpang kunyit (*Curcuma domestica* Val.) dan ekstrak kulit batang mimba (*Azadirachta indica* A. Juss) masing-masing menunjukkan aktivitas gastroprotektif baik pada uji preklinik maupun klinik. Tujuan dari penelitian ini adalah untuk mengetahui efektivitas gastroprotektif kombinasi ekstrak rimpang kunyit dan kulit batang mimba pada tikus yang diinduksi dengan asetosal.

Penelitian ini menggunakan tikus jantan galur Sprague Dawley dengan berat badan 100- 150 gram sebanyak 32 ekor yang dibagi menjadi 8 kelompok dengan metode Rancangan Acak Lengkap (RAL); kelompok I, II, III, IV, V, dan VI diberikan kombinasi ekstrak selama 7 hari sebelum diinduksi, kelompok VII sebagai kontrol negatif diberikan CMC 1% selama 7 hari sebelum dinduksi, kelompok VIII sebagai kontrol normal diberikan larutan CMC 1% dan tidak diinduksi dengan asetosal. Delapan jam setelah perlakuan, tikus dibedah dan dilakukan pengujian pada organ lambung meliputi perhitungan indeks ulkus, pemeriksaan keasaman lambung, determinasi mukus, dan pengamatan histologi. Hasil dari penelitian ini menyebutkan bahwa pemberian kombinasi ekstrak rimpang kunyit (50 mg/kg bb) dan ekstrak kulit batang mimba (250 mg/kg bb) dapat menurunkan indeks ulkus secara signifikan, pembentukan mukus yang tinggi, serta didukung dengan hasil pengamatan histologi, yakni tidak ditemukannya perubahan bentuk dan ukuran sel parietal yang bermakna jika dibandingkan dengan kontrol normal.

.....Some studies reported that turmeric (*Curcuma domestica* Val.) rhizome and neem (*Azadirachta indica* A. Juss) bark extracts which each of them demonstrated strong gastroprotective activities on both preclinical and clinical studies. The objective of this study was to investigate the gastroprotective effect of turmeric rhizome and neem bark extracts on acetosal-induced gastric mucosal lesions in rats. Thirty two male Sprague Dawley rats 100-150 g bw used in the study were divided into 8 groups using Complete Randomized Design (CRD) method; group I, II, III, IV, V and VI received combination extracts orally in various doses for 7 days before acetosal, group VII served as negative control received orally 1% CMC for 7 days before acetosal, group VIII received orally 1% CMC solution and served as normal control. Eight hours after treatment, animals were sacrificed and the stomach were taken to measure ulcer index, gastric acid determination, mucus determination and histology examination. The result suggests that pretreatment with combination of turmeric rhizome (50 mg/kg bw) and neem bark extract (250 mg/kg bw) and was observed significantly reduced the ulcer index, demonstrated high mucus production, also supported by histopathological examination with no significantly changing on parietal cells microscopic appearance compared with normal control.