

Ekstraksi annonaceous acetogenin dari daun sirsak, annona muricata, sebagai senyawa bioaktif antikanker

Monica Wijaya, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20289102&lokasi=lokal>

Abstrak

Senyawa bioaktif yang berasal dari tanaman sirsak atau *Annona muricata* telah lama diteliti dan terbukti bersifat antikanker, selain itu juga bersifat antiparasit, insektisida, anticacing, antibakteri, dan antivirus. Senyawa bioaktif, yang diekstrak dari daun sirsak, bernama annonaceous acetogenin dan telah berhasil dilakukan analisa secara kualitatif. Penelitian bertujuan untuk mendapatkan ekstrak yang kaya akan annonaceous acetogenin melalui maserasi daun sirsak yang sudah kering dengan pelarut etanol 95%, fraksinasi, dan kolom kromatografi. Senyawa annonaceous acetogenin yang terbukti larut dalam fraksi metanol 90% (Fraksi F005) dari tahapan fraksinasi dianalisis dengan LC-MS (Liquid Chromatography-Mass Spectroscopy) dengan adanya senyawa yang berat molekulnya 612 dan BST (Brine Shrimp Lethality Test) dengan nilai LC50 (Lethal Concentration) di bawah 1000 ppm. Dari kolom kromatografi terhadap fraksi F005 didapatkan 12 botol di antara 15 botol yang terbukti mengandung annonaceous acetogenin dari analisis TLC (Thin Layer Chromatography), LCMS, dan BST, dan selanjutnya dianalisis dengan HPLC untuk mengetahui banyaknya campuran senyawa yang terkandung di dalamnya.

.....Bioactive compound from soursop plants or *Annona muricata* has been investigated for a long time and proven to act as an anticancer, furthermore as parasiticidal, insecticidal, antibacterial, and antiviral. This bioactive compound, which is extracted from soursop leaves, is annonaceous acetogenin and succeeded in qualitative analysis. This aim of research is to get a rich content of annonaceous acetogenin by macerate the soursop leaves with ethanol 95%, fractinate, and isolate with column chromatography. Annonaceous acetogenin is proven to be soluble in methanol 90% fraction (Fraction F005) from fractination step which analyzed using LC-MS (Liquid Chromatography-Mass Spectroscopy) with 612 for its molecular weight and BST (Brine Shrimp Lethality Test) with LC50 below 1000 ppm. From column chromatography for fraction F005, it gave 12 bottles among all 15 bottles which are proven to contain annonaceous acetogenin by analyzing them using TLC (Thin Layer Chromatography), LC-MS, and BST, furthermore by HPLC to give knowledge about the mixture of compounds contained in them.