

## Kapasitas ekstrak Temulawak (*Curcuma xanthorrhiza* Roxb) sebagai anti *Streptococcus mutans* dalam menghambat demineralisasi email (in vitro)

Handoko Tirta, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20308067&lokasi=lokal>

---

### Abstrak

Xanthorrhizol yang di isolasi dari Temulawak dapat mempertahankan pH model biofilm in vitro selama 4 jam. Diketahui KBM ekstrak Temulawak terhadap *S. mutans* 25%.

Tujuan : Menganalisis efek ekstrak Temulawak 25% terhadap demineralisasi email yang terpapar biofilm *S. mutans*.

Metode : Model biofilm diperoleh dengan mengkultur *S. mutans* yang sudah ditanam dalam TYS broth selama 24 jam pada 6 well plate yang telah dilapisi pelikel. Ekstrak Temulawak dipaparkan pada model biofilm pada berbagai durasi antara 1-48 jam. Pengukuran pH menggunakan pH universal indicator. Model biofilm juga ditumbuhkan pada permukaan sample gigi. Pemaparan Ekstrak Temulawak dilakukan pada jam ke 16-20. Uji kekerasan mikro menggunakan indenter Knoop sebelum dan sesudah perlakuan.

Hasil : Sampai dengan jam ke 4, pH model biofilm yang terpapar Ekstrak Temulawak 25% tidak mengalami penurunan pH. Tidak terlihat efek Ekstrak Temulawak terhadap kekerasan permukaan email.

Kesimpulan : Ekstrak Temulawak 25% mampu menghambat penurunan pH biofilm, tetapi tidak berpengaruh terhadap demineralisasi email.

.....

Xanthorrhizol isolated from Java turmeric is able to maintain the pH of biofilm model in vitro for 4 hours. It was known that MBC of Java turmeric extract was 25%.

Purpose: To analyse the effect of 25% Java turmeric extract on email demineralization exposed to *S. mutans* biofilm.

Methods: Biofilm model was obtained by culturizing *S. mutans* which was cultured on TYS Broth during 24 hours on 6 well- plates which was layered by pellicle. Java turmeric Extract was added to biofilm model at various duration between 1-48 hours. pH measurement using pH universal indicator. Biofilm model was also cultured at tooth sample surface. Java turmeric extract was added at 16-20 hours. Micro hardness test was conducted using Knoop indenter before and after the intervention.

Result : After 4 hours, the pH of biofilm model which was exposed to Java turmeric 25% was not decreasing. No difference was found on the enamel micro hardness between experiment and control groups.

Conclusion : Java turmeric 25% is able to prevent reduction of biofilm pH, but does not have effect on enamel demineralization.