

Tingkat kebisingan lalu lintas dan risiko hipertensi pada supir angkutan umum KWK wilayah Jakarta Timur tahun 2012 = Road traffic noise and risk of hypertension on driver of public transportation in East Jakarta 2012

Vina Anggraeni, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20318268&lokasi=lokal>

Abstrak

Kebisingan lalu lintas merupakan salah satu masalah kesehatan lingkungan yang menimbulkan efek buruk bagi kesehatan pada orang yang terpapar, salah satu efeknya adalah hipertensi. Tujuan utama penelitian ini adalah untuk mengetahui hubungan antara tingkat kebisingan lalu lintas dengan risiko hipertensi.

Studi yang digunakan adalah cross sectional. 100 responden ditentukan menggunakan metode quota sampling pada supir angkutan umum KWK T.16, KWK T.06, dan KWK T.08. Pengukuran kebisingan menggunakan Sound Level Meter, dilakukan sebanyak 30 titik di sepanjang trayek yang dilalui responden dengan menggunakan purposive sampling sebagai penentuan titik sampling.

Prevalensi hipertensi diketahui melalui pengukuran langsung menggunakan sphygmomanometer air raksa dan stetoskop. Informasi mengenai faktor individu, faktor paparan, dan faktor perilaku juga diamati pada penelitian ini. Data yang didapat kemudian diolah menggunakan uji kai kuadrat dan regresi logistik.

Hasil studi menunjukkan tidak ada hubungan yang signifikan untuk tingkat kebisingan lalu lintas (Leq, 8hr) dengan risiko hipertensi ($p=0,135$). Hasil yang signifikan hanya didapatkan untuk faktor suku ($p=0,024$) dan obesitas ($p=0,049$). Tingginya prevalensi hipertensi pada penelitian ini tidak berhubungan dengan tingkat kebisingan lalu lintas tetapi berhubungan dengan suku dan obesitas. Oleh karena itu, bagi responden disarankan untuk menurunkan berat badan dengan melakukan olahraga secara rutin dan melakukan program diet.

.....Traffic noise is one of the environmental health problems that cause adverse health effects in people exposed, one effect is hypertension. The main purpose of this study was to determine the relationship between traffic noise levels with the risk of hypertension.

The study used a cross sectional. 100 respondents was determined using quota sampling method on drivers of public transport KWK T.16, KWK T.06, and KWK T.08. Noise measured using a Sound Level Meter, performed a total of 30 points along the route through which the respondents by using purposive sampling as the determination of the sampling point.

The prevalence of hypertension was known through direct measurement used a mercury sphygmomanometer and stethoscope. Information on individual factors, exposure factors, and behavioral factors were also observed in this study. Data was then processed using kai square test and logistic regression.

The study results showed no significant relationship to the traffic noise level (Leq, 8hr) with risk of hypertension ($p = 0.135$). The results obtained were significant only for ras factor ($p = 0.024$) and obesity ($p = 0.049$). The high prevalence of hypertension in this study did not correlate with levels of traffic noise but related to race and obesity. Therefore, the respondent advised to lose weight by exercising regularly and doing a diet program.