

Hubungan antara status gizi, asupan gizi, dan aktivitas fisik dengan VO₂max pada Mahasiswa Program Studi Gizi FKM UI tahun 2012 = Relationship between nutritional status, nutrient intake and physical activity with VO₂max in Undergraduate Students of Public Health University of Indonesia Majoring Nutrition in 2012

Eko Cipako Sinamo, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20319889&lokasi=lokal>

Abstrak

Skripsi ini membahas hubungan antara indeks massa tubuh (IMT), persen lemak tubuh (PLT), asupan zat gizi makro (kalori, karbohidrat, lemak dan protein), asupan zat gizi mikro (tiamin, riboflavin, piridoksin, vit.C dan Fe), dan aktivitas fisik dengan VO₂max. Penelitian ini adalah penelitian kuantitatif dengan desain cross sectional yang dilakukan pada 81 mahasiswa Reguler Gizi Kesehatan Masyarakat FKM UI angkatan 2010 dan 2011. VO₂max diukur dengan menggunakan alat Fitmate Med Hasil uji korelasi menunjukkan hubungan negatif antara IMT ($r = -0,231$) dan persen lemak tubuh ($r = -0,447$) dengan VO₂max pada responden keseluruhan. Terdapat hubungan positif antara asupan Fe ($r = 0,231$), dan aktivitas fisik ($r = 0,338$) dengan VO₂max pada responden keseluruhan. Diperlukan penelitian lebih lanjut dengan sampel yang lebih besar pada atlet dengan pengendalian yang lebih ketat terhadap faktor-faktor lain yang berpotensi menyebabkan bias dalam penelitian agar korelasi variabel independen dengan data VO₂max dapat merepresentasikan kekuatan hubungan yang sebenarnya.

.....This thesis discusses the relationship between body mass index (BMI), body fat percent (BFP), the intake of macro nutrients (calories, carbohydrates, fats and proteins), the intake of micro nutrients (thiamin, riboflavin, pyridoxine, vit. C and Fe), and physical activity with VO₂max. The study was a quantitative study with cross sectional design conducted in 81 undergraduate students of Public Health University of Indonesia majoring Nutrition in 2012. VO₂max was measured by using Fitmate Med. The result of correlation test showed a negative relationship between BMI ($r = -0,231$) and percent body fat ($r = -0,447$) with VO₂max in the overall respondents. Artifacts positive association between intake of Fe ($r = 0,231$) and physical activity ($r = 0,338$) with VO₂max in the overall respondents. There were no significant relationship between other independent variables with VO₂max. Further research is needed with larger samples in athletes with a more strict control of other factors that could potentially lead to bias in the study so that the data correlation with VO₂max independent variables can represent the real strength of the relationship.