

Isolasi dan penentuan struktur serta uji bioaktivitas senyawa kimia dari Daun Binahong Anredera cordifolia (Ten.) Steenis = Isolation, chemical structure elucidation and bioactivity test of the chemical compound of Binahong Leaves Anredera cordifolia (Ten.) Steenis

Ratna Djamil, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20329071&lokasi=lokal>

Abstrak

A research has been conducted to reveal the isolation and identification of bioactive constituents of leaves *Anredera cordifolia* (Ten.) Steenis (local name known as binahong) and obtained from the plantation of medicinal plants in BALITTRO, Lembang. Dried leaf samples were extracted with methanol and the results were fractionated with n-hexane, ethyl acetate and n-buthanol respectively. It was tested biological activity ,to toxicity on Brine shrimp test of *A.salina* Leach, an antioxidant with the DPPH method, antimicrobial and cytotoxicity test against murine P388 leukemia cells and breast cancer cells T-47D. Isolation was carried out by column chromatography techniques, followed by purification using preparative thin layer chromatography techniques and preparative HPLC. Determination of molecular structure performed by analyzing the UV-Vis spectrum data, Infrared, LC-MS, 1H-NMR, 13C-NMR, DEPT, 2D-NMR include HMQC and HMBC. The results of isolation of the leaf extract of *A.cordifolia* were obtained three compounds, a compound that is proposed as a new compound in this plant that is 8-Glucopyranosyl-4?,5,7-trihydroxyflavone and two other compounds are adenine and (9?Z,?Z)-propane-1,2,3-tryl trioleat. The results of bioactivity BS LT, methanol extract, n-hexane, ethyl acetate, n-buthanol and isolates 8-Glucopyranosyl-4?,5,7-trihydroxyflavone were toxic to shrimp larvae *A.salina* with LC50 values : 53,11; 256,23; 57,96; 132,39; and 68,07μg/mL. The results antioxidant activity, all of extract active except for n-hexane fraction with IC50 values 53,11; 256,23; 57,96; 132,39 and 68,07 μg/mL. The results of the antibacterial activity against *S.aureus*, *E.coli* and *C.albicans* , n-hexane extracts had no inhibitory power, the ethyl acetate extract was more active than the methanol and n-buthanol extract against bacteria *S.aureus* at a concentration of 500 mg/1 mL. Methanol extract of leaves of *A.cordifolia* was not potentially active against murine P388 and T-47D cells, ethyl acetate extract was shown activity against murine P388 cells with IC50 values 62,74 μg/mL. The compounds 8-Glucopyranosyl-4?,5,7-trihydroxyflavone, not active against T-47D cells but was active against P388 cells with IC50 values 87,13 μg/mL. Isolate adenine was active against murine P388 cells and breast cancer cells T-47D with IC50values 89,08 and 39 μg/mL, respectively.

<hr>Telah dilakukan penelitian isolasi dan identifikasi untuk mengungkapkan kandungan senyawa kimia dari daun tanaman *Anredera cordifolia* (Ten.) Steenis yang dikenal dengan nama binahong dan diperoleh dari perkebunan tanaman obat BALITTRO , Lembang. Sampel daun kering diekstraksi dengan metanol dan hasilnya difraksinasi dengan n-heksana, etil asetat dan n-butanol. Setiap fraksi dilakukan uji aktivitas, yang meliputi uji toksisitas terhadap larva udang *A. salina* Leach, antioksidan dengan metode DPPH, antimikroba serta uji sitotoksitas terhadap sel leukemia murine P388 dan sel kanker payudara T-47D. Isolasi dilakukan dengan teknik kromatografi kolom, dilanjutkan dengan pemurnian menggunakan teknik kromatografi lapis tipis preparatif dan HPLC preparatif. Penentuan struktur molekul dilakukan dengan menganalisis data spectrum UV-Vis, Infra merah, LC-MS, 1H-NMR, 13C-NMR, DEPT, 2D-NMR meliputi

HMQC dan HMBC . Hasil isolasi terhadap ekstrak daun *A. cordifolia* diperoleh 3 senyawa, yaitu satu senyawa yang diusulkan sebagai senyawa baru pada tanaman *A.cordifolia* yaitu 8-Glucopyranosyl-4',5,7-trihydroxyflavone, dan 2 senyawa lainnya yaitu senyawa adenine dan senyawa (9'Z,9''Z)-propane-1,2,3-triyl trioleat. Berdasarkan hasil uji bioaktivitas BSLT, ekstrak metanol, n-heksana, etil asetat, n-butanol dan isolat 8-Glucopyranosyl-4',5,7-trihydroxyflavone toksik terhadap larva udang *A. salina* masing-masing mempunyai nilai LC₅₀ 46,19; 542,05; 32,06; 79,72 dan 24,74 μg/mL. Hasil uji aktivitas antioksidan seluruh ekstrak bersifat aktif, kecuali ekstrak n-heksana tidak aktif, masing-masing dengan nilai IC₅₀ 53,11, 256,23 57,96 , 132,39 dan 68,07μg/mL. Hasil uji aktivitas antibakteri terhadap *S. aureus*, *E.coli* dan *C. albicans*, ekstrak n-heksana tidak memiliki daya hambat, ekstrak etil asetat lebih aktif dibandingkan dengan ekstrak metanol, dan n-butanol terhadap bakteri *S.aureus*, pada konsentrasi 500 mg/1mL. Ekstrak metanol daun *A. cordifolia* tidak mempunyai potensi aktif terhadap sel murine P388 dan T-47D, tetapi ekstrak etil asetat mempunyai potensi aktif terhadap sel murine P388 dengan IC₅₀ 62,74 μg/mL. Senyawa 8-Glucopyranosyl-4',5,7-trihydroxyflavon, tidak aktif terhadap sel T-47D tetapi aktif terhadap sel P388 dengan nilai IC₅₀ 87,13 μg/mL. Isolat adenine mempunyai potensi aktif terhadap sel murine P388 maupun sel kanker payudara T-47D dengan nilai IC₅₀ 89,08 dan 39 μg/mL.