

Profil keberadaan Candida spp pada Vagina ibu hamil dengan persalinan Preterm dan Neonatus yang dilahirkannya = Profile of Candida spp in the Vagina of pregnant women with Preterm labor and their Neonates / Ruswanto Hadi Sucipto

Ruswanto Hadi Sucipto, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20330264&lokasi=lokal>

Abstrak

<ABSTRAK

Keberadaan Candida pada vagina ibu hamil dapat menjadi sumber infeksi bagi bayi yang dilahirkannya saat melalui jalan lahir atau selama masa perawatan bayi. Penelitian ini bertujuan mengetahui profil Candida spp pada vagina ibu hamil yang mengalami persalinan preterm dan neonatus yang dilahirkannya. Penelitian berdesain potong lintang ini menggunakan teknik consecutive sampling pada ibu hamil trimester III dengan persalinan preterm yang datang ke RSCM serta neonatus yang dilahirkannya. Terdapat 38 ibu hamil dengan persalinan preterm yang mengikuti penelitian sehingga diperoleh 38 bahan klinik cairan vagina ibu, serta 22 usap mulut dan 10 usap perianal neonatus. Pada bahan klinik tersebut dilakukan pemeriksaan mikologi untuk mengetahui frekuensi dan spektrum spesies Candida. Hasil penelitian menunjukkan frekuensi keberadaan Candida spp pada vagina ibu hamil yang mengalami persalinan preterm sebesar 47,4 %. Profil Candida spp dari cairan vagina ibu terdiri atas: Candida albicans 12 isolat (66,7%), Candida tropicalis 3 isolat (16,7%), Candida glabrata 2 isolat (11,1%) dan Candida parapsilopsis 1 isolat (5,5%). Frekuensi keberadaan Candida pada usap mulut neonatus sebesar 13,6% sedangkan pada usap perianal ditemukan Candida pada 2 dari 10 neonatus. Profil Candida spp pada neonatus terdiri atas: Candida albicans dan Candida parapsilopsis. Hubungan antara keberadaan Candida spp pada vagina ibu hamil yang mengalami persalinan preterm dengan Candida spp pada neonatus belum dapat diketahui karena keterbatasan penelitian.

<hr>

ABSTRACT

The presence of Candida in vagina of pregnant women can be a source of infection in their neonates when pass birth canal or during baby care. The aim of this study is to determine the profile of Candida spp in vagina of pregnant women with preterm labor and their neonates. This cross-sectional study was conducted with consecutive sampling techniques on third-trimester pregnant women with preterm labor who come to Cipto Mangunkusumo hospital and their neonates. There were 38 pregnant women with preterm labor, and obtained 38 vaginal swab of the mothers, 22 oral swab and 10 perianal swab of their neonates. Mycological examination was

conducted to determine the frequency and spectrum of *Candida* species. The results showed frequency of *Candida* spp in the vagina of pregnant women with preterm labor was 47.4%. The spectrum of *Candida* spp from 38 vaginal swab of the mothers consists of *Candida albicans* 12 isolates (66.7%), *Candida tropicalis* 3 isolates (16.7%), *Candida glabrata* 2 isolates (11.1%), and *Candida parapsilopsis* 1 isolate (5.5%). The frequency of *Candida* spp of 22 oral swab was 13.6% and 2 of 10 of perianal swab. The spectrum of *Candida* spp in neonates consists of *Candida albicans* and *Candida parapsilopsis*. The relationship between the presence of *Candida* spp in vagina of the mothers and their neonates could not be known yet due to the limitation of this study.