

Mendeteksi kinetik antigen ns 1 dengan menggunakan strip bio rad ns 1 ag di pasien yang terdeteksi dengue pada hari demam ke 4 6 di indonesia = Determining the kinetic of ns 1 antigen using bio rad ns 1 ag strip in dengue infected patients at day 4 to 6 of fever in indonesia

Emawulidasari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20348329&lokasi=lokal>

Abstrak

Protein Non Structural 1 NS 1 telah digagaskan untuk menjadi alat diagnostik dini untuk infeksi demam berdarah karena keberadaannya yang cukup tinggi pada saat fase akut dari infeksi ini NS 1 adalah protein yang sangat penting bagi virus dengue untuk melakukan replikasi virus Penelitian ini bertujuan untuk menetapkan kinetik NS 1 pada hari demam ke 4 sampai dengan 6 dengan menggunakan Strip Bio Rad Ag di komunitas di Jakarta Seluruh penelitian ini berjalan selama 36 bulan Maret 2010 ndash Februari 2013 Terdapat 102 pasien diduga terinfeksi demam berdarah yang memenuhi kriteria inklusi Serum darah pasien dites menggunakan RT PCR atau isolasi virus di line sel C6 36 atau kenaikan titer antibodi sebagai standar baku dan sedangksn Strip Bio Rad NS 1 Ag digunakan sebagai metode diagnostik yang digunakan dalam penelitian ini SPSS 17 0 merupakan alat statistik yang digunakan dalam riset ini Dari kesuluruhan 68 68 3 pasien terbukti positive terinfeksi dan 34 31 7 merupakan negatif untuk infeksi dengue Sementara itu hanya 54 52 94 pasien yang terdiagnosis positif infeksi dengue pada hari demam ke 4 6 Kinetik NS 1 selama hari demam ke 4 6 terdeteksi masing masing sebanyak 83 64 69 09 and 47 27 Kinetik NS 1 pada pasien dengue infeksi primer sejak hari demam ke 4 sampai dengan 6 terdeteksi sebanyak 100 83 33 dan 75 Secara kontras kinetik NS 1 di pasien degue infeksi sekunder dari hari demam ke 4 6 adalah 73 33 60 dan 26 67 Terdeteksinya NS 1 akan semakin menurun seiring dengan bertambahnya hari demam Antingen NS 1 pada pasien dengue infeksi primer lebih lama keberadaannya dibandingkan dengan pasien dengue infeksi sekunder.

.....NS 1 has been purposed to be use as an early method to diagnose dengue infection because of its appearance in the acute phase of the infection NS 1 is a protein that essential for dengue virus to do the virus replication This study has aimed to determining the kinetic of NS 1 by using Bio Rad NS 1 Ag Strip for dengue infection during day 4 6th of fever in community in Jakarta The entire study was conducted in 36 months March 2010 ndash February 2013 There were 102 suspected dengue infection patients that fulfill the inclusion criteria tested for the NS 1 antigen in their blood serum using RT PCR or isolation of the viruses in C6 36 cell line or increasing antibody titer as the gold standard Meanwhile Bio Rad NS 1 Ag Strip was used as the diagnostic method use in this study Data were analyzed using SPSS 17 0 Overall 68 68 3 patients were considered positive and 34 31 7 patients were negative for dengue infection Meanwhile only 54 52 94 patients were positive at the day 4 6th of fever The kinetic of NS 1 during day 4 6th day of fever were 83 64 69 09 and 47 27 respectively The kinetic of NS 1 in primary patients from day 4 to day 6 were 100 83 33 and 75 respectively In contrast the kinetic of NS 1 in secondary patients from day 4 to day 6 respectively were 73 33 60 and 26 67 The presences of NS 1 antigen were decreasing as the day of fever keeps progressing The availability of NS 1 antigen in primary dengue infected patients was longer than the secondary dengue infected patients.