

Faktor-faktor yang berhubungan dengan praktik kewaspadaan universal oleh perawat rumah sakit = The factors associated with the practice of universal precautions by the hospital's nurse / Eka Syarifah Dewi

Eka Syarifah Dewi, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20349943&lokasi=lokal>

Abstrak

**ABSTRAK
**

Latar Belakang: Perawat mempunyai risiko terinfeksi virus yang ditularkan melalui darah seperti virus HIV/AIDS, Hepatitis B dan Hepatitis C. Penularan dapat dicegah dengan penerapan prinsip kewaspadaan universal. Tujuan : Diketahuinya gambaran praktik kewaspadaan universal oleh perawat di RSUD Karawang dan hubungannya dengan faktor predisposisi, faktor pemungkin dan faktor penguat. Metode Penelitian: Penelitian deskriptif analitik dengan metode potong lintang yang dilaksanakan pada bulan Juni-Juli 2013. Data diperoleh dari 77 responden melalui self administered questionnaires dan observasi. Hasil : Penelitian menunjukkan hanya 44,2% responden yang melakukan praktik kewaspadaan universal dengan baik. Dari hasil uji chi square, variabel yang secara signifikan berhubungan dengan praktik kewaspadaan universal adalah pengetahuan ($p = 0,045$) dan pengawasan atasan ($p = 0,011$) serta dukungan teman sekerja ($p = 0,037$). Variabel yang tidak berhubungan adalah pendidikan, lama masa kerja, sikap, ketersediaan SOP, ketersediaan fasilitas dan pelatihan.

Kesimpulan : Faktor yang paling berhubungan dengan praktik kewaspadaan universal adalah pengawasan atasan. Responden yang mendapatkan pengawasan atasan mempunyai peluang lebih dari 3 kali untuk melakukan praktik kewaspadaan universal dibandingkan yang tidak mendapatkan pengawasan atasan.

<hr>

**ABSTRACT
**

Background : The nurse has a risk to be infected by the virus that is infected blood borne pathogens through the blood such as the virus of HIV/AIDS, Hepatitis B and Hepatitis C. The infection can be prevented by the principle of Universal Precautions. Objective: Knowledgeable overview of the practice of universal precautions by nurses in RSUD Karawang and the associated with predisposition factor, enabling factors and reinforcing factors. Methods: The research uses the analytic descriptive with a crosssectional method which is held on June-July 2013. The data is obtained from 77 respondents through self administered questionnaires and observation. Results: Result of research shows only 44.2% of respondents who practice universal precautions as well. From the results of the chi square test, the variables that were significantly associated with the practice of universal precautions is knowledge ($p = 0.045$) and control superiors ($p = 0.011$) as well as the support of co-workers ($p = 0.037$). Variables are unrelated is education, length of service, attitude, availability of SOP, availability of facilities and training. Conclusions: Factors most associated with the

practice of universal precautions is superior supervision. Respondents who have a chance of getting control superiors more than 3 times to practice universal precautions than not getting control superiors.