

Risiko tingkat pajanan pestisida terhadap penyakit asma pada petani di Kabupaten Karo tahun 2012 = The risk of level exposure pesticide for incident asthma in farmer at Karo District in 2012

Munthe, Simeon, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20350308&lokasi=lokal>

Abstrak

Terpapajanan pestisida secara kronis dapat mempengaruhi status kesehatan. Efek tersebut bergantung pada toksisitas pestisida dan tingkat pajanannya. Bisnis petani kebanyakan memerlukan pestisida untuk meningkatkan hasil pertanian. Petani yang menggunakan pestisida harus memakai APD untuk mengurangi dampak yang ditimbulkan oleh pestisida.

Desain kasus-kontrol telah digunakan untuk mengetahui risiko pajanan pestisida pada asma di kalangan petani, dengan 70 sampel yang menderita asma dan 210 sampel kontrol yang tidak asma, bertempat tinggal di desa dan bekerja sebagai petani, telah dilakukan di Kabupaten Karo Provinsi Sumatera Utara. Variabel confounding dalam penelitian ini adalah penggunaan APD dan variabel tingkat pendidikan. Dalam penelitian ini tingkat eksposur dan penggunaan APD dikategorikan menjadi tiga kategori.

Hasil analisis: Risiko kejadian asma pada tingkat pajanan sedang dibandingkan dengan rendah (OR = 2,33, 95% CI: 0,72 – 7,61) sedangkan risiko kejadian asma pada tingkat pajanan tinggi dibandingkan dengan rendah (OR = 3,24, 95% CI: 1,06 – 10,37). Risiko kejadian asma pada penggunaan APD sedang dibandingkan dengan kurang (OR = 0,37, 95% CI: 0,19 – 0,72), sedangkan risiko kejadian asma pada penggunaan APD baik dibandingkan dengan penggunaan APD buruk (OR = 0,2, 95% CI: 0,07 – 0,53). Efek tingkat pajanan terhadap kejadian asma dipengaruhi oleh penggunaan APD, semakin lengkap penggunaan APD semakin kecil efek tingkat pajanan terhadap kejadian asma.

.....

Chronic exposed of pesticide have harmful effect on health. Those harmful effect depends on the level of exposure and toxicity of pesticide. The farmer bussiness mostly need pesticide achieve the optimal of agriculture results. The farmer who use pesticide must be use personal protected equipments to reduced harmful effect of pesticide.

Design case-control have been use to study the risk of pesticide exposure on asthma among farmers, with 70 sampel who suffer from asthma and 210 control sampel who are not asthmatic, residing in the village and worked as farmer, has been done in Karo district of North Sumatera Province. The confounding in this study are use PPE variable and education level variable. In this study the level of exposure and use PPE are categorized into three categories.

Result analysis: the effect of middle level exposure of pesticide compared to low level on asthma (OR = 2.33, 95% CI: 0.72 to 7.61), while the effect of high level compared with low exposure on asthma (OR = 3.24, 95% CI: 1.06 to 10.37). the effect of middle level usage of PPE compared to less on asthma (OR = 0.37, 95% CI: 0.19 to 0.72), while the effect of good usage of PPE compared to less (OR = 0.2, 95% CI : .07 to .53). Effects of exposure level of pesticide on asthma is reduced by the use of PPE, more complete usage of PPE more diminishing the effect of exposure level on asthma.