

Relasi antara infeksi trichuris trichiura dengan status anemia pada anak-anak yang tinggal di Kabupaten Nangapanda Nusa Tenggara Timur = The Relationship between trichuris trichiura infection and anemia status of the children living in Nangapanda subdistrict East Nusa Tenggara

Jason Rahmadi Ruslie, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20385599&lokasi=lokal>

Abstrak

Trichuris trichiura adalah soil-transmitted helminths (STH) yang umum ditemukan di seluruh dunia, terutama di daerah tropis dengan sanitasi yang buruk. Banyak anak-anak usia prasekolah dan sekolah tinggal di daerah dimana parasit ini secara intensif ditransmisikan, dan membutuhkan pengobatan dan intervensi pencegahan segera. Tujuan dari penelitian ini adalah untuk menentukan prevalensi anemia pada anak-anak baik yang terinfeksi *T. Trichiura* maupun anak-anak yang tidak terinfeksi yang tinggal di daerah endemik desa Nangapanda, Kabupaten Ende, Nusa Tenggara Timur. Penelitian ini dilakukan dari bulan Mei sampai Juni 2010. Metode analisa feces yang digunakan adalah konsentrasi formol-ethyl asetat and analisa darah dengan menggunakan alat sysmex KX 21 untuk mengukur anemia. Penelitian ini menggunakan desain cross sectional analitik. Data sekunder diperoleh dari Departemen Parositologi, Fakultas Kedokteran Universitas Indonesia. Jumlah peserta bertotal 262 anak.

Penelitian ini menunjukkan bahwa risiko terinfeksi *Trichuris* pada anak usia 12-14,99 tahun secara signifikan lebih rendah dibandingkan dengan anak-anak dalam kelompok usia 5-11,99. Namun, korelasi antara infeksi *T. trichiura* dan status anemia ditemukan tidak signifikan bahkan setelah disesuaikan dengan jenis kelamin dan usia anak. Pada kesimpulan, tidak ada hubungan yang signifikan antara infeksi *trichiura* *Trichuris* dengan status anemia peserta penelitian. Penelitian lebih lanjut dibutuhkan untuk memperjelas adanya hubungan *Trichuris trichiura* dan anemia, beserta dengan jenis cacing lainnya.

.....*Trichuris trichiura* is a soil-transmitted helminth (STH) which is commonly found throughout the world, especially in tropical areas with poor sanitation. Many preschool-age children and school-age children live in areas where these parasites are intensively transmitted. The objective of this study was to determine the prevalence of anemia in *T. trichiura* infected children and non-infected children living in endemic area of Nangapanda village, Ende district, East Nusa Tenggara. The research was performed from May to June 2010. Formol-ethyl acetate concentration method was used to analyze the stool sample and blood analysis sysmex KX21 was used to measure anemia. This study used analytical cross sectional design. Secondary data was obtained from the Department of Parasitology, Faculty of Medicine University of Indonesia. Total participants were 262 children.

The risk of having *Trichuris* infection in children aged 12-14.99 years was significantly lower compared to those in the 5-11,99 age group. However, the relationship between *T. trichiura* infection and anemia status was not significant even after adjusted to age and gender. In conclusion, there were no significant relationship between *T. trichiura* infection with the anemia status of the participants. Further study by using cohort design should made to elucidate the relationship between *Trichuris trichiura* and anemia, including other types of helminthes as well.