

Hubungan ketersediaan sarapan dan faktor lainnya dengan pola sarapan Siswa/i SMA terpilih di Kabupaten Tangerang Tahun 2014 = The relationship between availability and other factors with breakfast pattern at selected Highschool Student in Tangerang 2014

Imas Arumsari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20386471&lokasi=lokal>

Abstrak

Kebiasaan melewatkan sarapan memperbesar risiko terjadinya obesitas, pola hidup sehat, dan penurunan kecerdasan, khususnya pada pelajar. Terlebih, remaja sedang mengalami proses growth spurt dimana terjadi peningkatan kebutuhan gizi. Kabupaten Tangerang memiliki tingkat kelulusan UN SMA terendah di provinsi Banten pada tahun ajaran 2013 lalu, sehingga perlu diteliti lebih lanjut mengenai pola sarapan remaja di wilayah ini.

Penelitian ini bertujuan untuk mengetahui hubungan antara ketersediaan sarapan dan faktor lainnya dengan pola sarapan siswa/i SMA terpilih di Kabupaten Tangerang tahun 2014. Penelitian ini merupakan penelitian kuantitatif dengan desain studi cross sectional. Data dikumpulkan menggunakan kuesioner isian mandiri yang diisi 160 siswa. Asupan sarapan dinilai menggunakan kuesioner food recall 2x24 jam.

Hasil penelitian menunjukkan, sebanyak 71,8% responden memiliki pola sarapan buruk. Terdapat hubungan yang signifikan antara ketersediaan sarapan ($P=0,014$) dengan pola sarapan responden.

<hr>

Skipping breakfast habit increase the risk of obesity, unhealthy lifestyle, and lack of intelligence, especially the students. Moreover, adolescent growth spurt is undergoing a process whereby an increase in nutritional requirements. Tangerang has lowest high school graduation rate in Banten Province for the academic year 2013 so it is important to develop research on adolescent breakfast pattern in this region.

This study aims to determine the relationship between the availability of breakfast and other factors with student breakfast pattern in selected high school in Tangerang on 2014. Research method is a quantitative research with cross sectional design study. Data were collected using self-completed questionnaires for 160 students. Breakfast intake was assessed using a 2x24 hour food recall questionnaire.

The results showed, 71,8% of respondents had a bad breakfast pattern. There is a significant association between the availability of breakfast ($P=0.014$) with respondents' breakfast pattern.