

Studi pembentukan dna adduct 8 hidroksi 2 deoksiguanosin 8 ohdg dari senyawa propil galat pg dan 2 6 di tert butil p benzoquinon bht quinon terhadap calf thymus dna dan 2 deoksiguanosin yang dimediasi oleh cupri klorida secara in vitro = In vitro study of formation of 8 hydroxy 2 deoxyguanosine 8 ohdg in calf thymus dna and 2 deoxyguanosine treated with propyl gallate pg and 2 6 di tert butyl p benzoquinone bht quinone mediated by cupric chloride

Dwi Retno Widiastuti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20389503&lokasi=lokal>

Abstrak

[ABSTRAK

Kerusakan oksidatif DNA yang disebabkan oleh propil galat (PG) dan 2,6-di-tertbutil-p-benzoquinon (BHT-Quinon, metabolit BHT), dianalisis dari pembentukan DNA adduct, 8-hidroksi-8-OHdG, terhadap Calf thymus DNA dan DNA base, 2-deoksiguanosin (dG). PG dengan dimediasi oleh CuCl₂ menyebabkan peningkatan 8-OHdG terhadap Calf thymus DNA sebesar 9,17 kali lebih besar dibandingkan terhadap kontrol (DNA tanpa perlakuan). Dengan adanya CuCl₂ pada konsentrasi 1,28.10⁻⁵ M, rasio pembentukan 8-OHdG dari hasil interaksi antara dG dengan PG pada berbagai variasi konsentrasi (20 - 150 ppm) berkisar antara 75,50 - 312,06 8-OHdG terhadap 105 dG. Pembentukan 8-OHdG tersebut, meningkat dengan bertambahnya konsentrasi PG dari 20 - 80 ppm, kemudian mulai menurun dengan bertambahnya konsentrasi PG. BHT-quinon, dengan adanya CuCl₂ menyebabkan peningkatan 8-OHdG terhadap Calf thymus DNA sebesar 0,05 kali dibandingkan kontrol (DNA tanpa perlakuan). Analisis menggunakan LC-MS/MS dilakukan untuk mengidentifikasi 8-OHdG, dengan puncak induk (M⁺. + 1) 284 dan memiliki dua fragmen utama m/z 167,9 dan m/z 139,9.

<hr>

ABSTRACT

Oxidative DNA damage caused by propyl gallate (PG) and 2,6-di-tert-butyl-pbenzoquinone (BHT-Quinone, a metabolite of butylated hydroxytoluene (BHT)), was evaluated by measuring the formation of DNA adduct, 8-hydroxy-2-deoxyguanosine (8-OHdG), in Calf thymus DNA and DNA base, 2-deoxyguanosine (dG). PG mediated with CuCl₂ increased 8-OHdG formation in Calf thymus DNA 9.17 fold from control (DNA without treatment). In the present of CuCl₂ 1.28.10⁻⁵ M, ratio 8-OHdG resulted from interaction of dG with PG at various concentration (20 - 150 ppm), was ranged from 75.50 - 312.06 8-OHdG

per 105 dG. This formation was increased by PG in a concentration-dependent manner ranged from 20 ppm up to 80 ppm, then decreased upon increasing the PG concentration. Meanwhile, BHT-quinone increased 0.05 fold from control (DNA without treatment) in the presence of CuCl₂. LC-MS/MS analysis was performed to identify molecular structure of 8-OHdG, which had base peak (M⁺. + 1) 284 and had two main fragment at m/z 167.9 and m/z 139.9., Oxidative DNA damage caused by propyl gallate (PG) and 2,6-di-tert-butyl-pbenzoquinone (BHT-Quinone, a metabolite of butylated hydroxytoluene 􀂱 BHT), was evaluated by measuring the formation of DNA adduct, 8-hydroxy-􀀕􀂶-deoxyguanosine (8-OHdG), in Calf thymus DNA and DNA base, 􀀕􀂶-deoxyguanosine (dG). PG mediated with CuCl₂ increased 8-OHdG formation in Calf thymus DNA 9.17 fold from control (DNA without treatment). In the present of CuCl₂ 1.28.10⁻⁵ M, ratio 8-OHdG resulted from interaction of dG with PG at various concentration (20 􀂱 150 ppm), was ranged from 75.50 􀂱 312.06 8-OHdG per 105 dG. This formation was increased by PG in a concentration-dependent manner ranged from 20 ppm up to 80 ppm, then decreased upon increasing the PG concentration. Meanwhile, BHT-quinone increased 0.05 fold from control (DNA without treatment) in the presence of CuCl₂. LC-MS/MS analysis was performed to identify molecular structure of 8-OHdG, which had base peak (M⁺. + 1) 284 and had two main fragment at m/z 167.9 and m/z 139.9.]