

# Faktor-faktor yang berhubungan dengan in-stent restenosis pada pasien pasca intervensi koroner perkutan = Factors related to in-stent restenosis in patients after percutaneous coronary intervention

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20391325&lokasi=lokal>

---

## Abstrak

[Latar Belakang. Angka kejadian In-Stent Restenosis (ISR) pasca Intervensi Koroner Perkutan (IKP) baik pada penggunaan Bare-Metal Stent (BMS) maupun Drug-Eluting Stent (DES) masih tinggi. Tujuan. Mengetahui faktor-faktor yang berhubungan dengan ISR pada pasien pasca IKP. Metode. Desain penelitian potong lintang retrospektif ini dilakukan dengan menggunakan rekam medik pasien pasca IKP yang menjalani follow-up angiografi di Pelayanan Jantung Terpadu/Rumah Sakit Umum Pusat Negeri Dr. Cipto Mangunkusumo dalam kurun waktu bulan Januari 2009 sd. Maret 2014. Gambaran angiografi ISR bila diameter stenosis pada follow-up angiografi lebih dan sama dengan 50 persen baik di dalam stent maupun menjulur keluar lima mm baik dari ujung proksimal maupun distal stent. Analisis multivariat pada penelitian ini dilakukan dengan menggunakan regresi logistik ganda. Hasil. 289 subyek penelitian terdiri dari 133 pasien dengan ISR dan 156 pasien tanpa ISR. Angka kejadian ISR pada penggunaan BMS dan DES masing-masing sebesar 61,3% dan 40,7%. Jenis stent (OR=4,83; 95% IK 2,51-9,30; p=0,001), panjang stent (OR=3,71; 95% IK 1,99-6,90; p=0,001), lesi di bifurkasi (OR=2,43; 95% IK 1,16-5,10; p=0,019), merokok (OR=2,30; 95% IK 1,33-3,99; p=0,003), diameter pembuluh darah (OR=2,18; 95% IK 1,2-3,73; p=0,005), hipertensi (OR=2,16; 95% IK 1,16-4,04; p=0,016) dan Diabetes Melitus/DM (OR=2,14; 95% IK; p=0,007) sebagai faktor prediksi ISR. Kesimpulan. Jenis stent, panjang stent, lesi di bifurkasi, merokok, diameter pembuluh darah, hipertensi dan DM merupakan faktor-faktor yang berhubungan dengan ISR pada pasien pasca IKP., Background. The incidence of In-Stent Restenosis (ISR) after Percutaneous

Coronary Intervention (PCI) both in the use of Bare-Metal Stent (BMS) and Drug-Eluting Stents (DES) are still high. Purpose. To determine factors related to ISR in patients after PCI. Method. A retrospective cross-sectional study was conducted using medical records of patients after PCI who underwent follow-up of angiography in the period between January 2009 to March 2014 in The Integrated Cardiac Service/Public Hospital Center Dr. Cipto Mangunkusumo Jakarta. Angiographic ISR was defined as diameter stenosis > 50% at follow-up angiography in the within of the stent, and within its five mm proximal and distal edges. Multivariate analysis performed in this study using regression multiple logistic. Results. 289 study subjects consisted of 133 patients with and 156 patients without ISR. The incidence of ISR in the use of BMS and DES,

respectively 61,3% and 40,7%. Using multivariate analysis, type of stent (OR=4,83; 95% CI 2,51-9,30; p=0,001), stent length (OR=3,71; 95% CI 1,99-6,90; p=0,001), bifurcation lesions (OR=2,43; 95% CI 1,16-5,10; p=0,019), smoking (OR=2,30; 95% CI 1,33-3,99; p=0,003), blood vessel diameter (OR=2,18; 95% CI 1,2-3,73; p=0,005), hypertension (OR=2,16; 95% CI 1,16-4,04; p=0,016) and Diabetes Mellitus/DM (OR=2,14; 95% CI; p=0,007) were identified as predictors of ISR. Conclusion. Type of stent, stent length, bifurcation lesions, smoking, blood vessel diameter, hypertension and DM were factors related to ISR in patients after PCI.]