

Pengaruh serotype c streptococcus mutans terhadap tingkat keasaman plak dan saliva pada penderita resesi gingiva dengan hipersensitif dentin (Analisis menggunakan real time PCR) = Effect of serotype c streptococcus mutans for plaque and saliva acidity levels in gingival recession with dentin hypersensitivity (Analysis using real time PCR)

Albert, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20403996&lokasi=lokal>

Abstrak

Latar Belakang: Hipersensitif dentin dipengaruhi oleh akumulasi plak pada permukaan gigi dan penetrasi bakteri pada tubulus dentin.

Tujuan: Menganalisis proporsi serotype c Streptococcus mutans dan tingkat keasaman di dalam plak dan saliva penderita resesi gingiva yang hipersensitif dentin dengan penderita resesi gingiva yang non hipersensitif.

Metode: Tiga puluh enam sampel plak dan saliva dibagi dalam dua kelompok yaitu kelompok hipersensitif dan non hipersensitif. Dilakukan ekstrak DNA sampel, pengukuran pH sampel dan evaluasi amplifikasi serotype c Streptococcus mutans dengan alat Real Time PCR.

Hasil: Proporsi serotype c Streptococcus mutans di dalam plak dan saliva tidak berbeda pada penderita resesi gingiva dengan hipersensitif dentin maupun non hipersensitif,

Kesimpulan: Proporsi serotype c Streptococcus mutans plak dan saliva tidak mempengaruhi hipersensitif dentin.

.....

Background: dentin hypersensitivity is affected by the accumulation of plaque on the tooth surface and penetration of bacteria in the dentinal tubules.

Objective: To analyze the proportion of serotype c Streptococcus mutans and the level of acidity in plaque and saliva of patients with hypersensitive dentin and non hypersensitive.

Methods: Thirty-six plaque and saliva samples were divided into two groups: the hypersensitive and non-hypersensitive. Extract the sample DNA, measure the acidity levels and evaluate serotype c Streptococcus mutans amplification with Real Time PCR.

Results: The proportion of serotype c Streptococcus mutans in plaque and saliva is not significantly different in the patients with gingival recession both hypersensitive and non-hypersensitive.

Conclusions: The proportion of serotype c Streptococcus mutans in plaque and saliva are equally well both in hypersensitive and non hypersensitive cases.