

Perbandingan ekspresi tissue inhibitor of metalloproteinase-2 (TIMP 2) pada ameloblastoma pleksiform dan folikuler secara imunohistokimia (Penelitian pada pasien ameloblastoma di Rumah Sakit Cipto Mangunkusumo periode Januari 2011-Agustus 2014) = Comparison of expression of tissue inhibitor of metalloproteinase-2 (TIMP 2) immunohistochemically in plexiform and follicular ameloblastoma (Research of ameloblastoma's patient in Rumah Sakit Cipto Mangunkusumo on January 2011-August 2014)

Muhammad Dhani Saleh, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20404428&lokasi=lokal>

Abstrak

Latar Belakang: Ekspresi TIMP-2 (Tissue Inhibitor of Metalloproteinase-2) secara imunohistokimia digunakan untuk menentukan sifat invasif lokal ameloblastoma yang berkaitan dengan kemampuan rekurensi.

Tujuan: Mengevaluasi ekspresi TIMP-2 secara imunohistokimia pada ameloblastoma pleksiform dan folikuler.

Metode: Dilakukan pemeriksaan imunohistokimia pada sampel ameloblastoma pleksiform ($n=16$) dan folikuler ($n=14$) dengan antibodi monoklonal TIMP-2. Ekspresi imunohistokimia TIMP-2 dinilai dengan software Image J.

Hasil: Tidak terdapat perbedaan bermakna ekspresi TIMP-2 pada ameloblastoma pleksiform dan folikuler pada uji Chi-Square dengan nilai signifikan $p=0.072$ ($p>0.05$).

Pembahasan: Ekspresi TIMP-2 yang lemah berkaitan dengan meningkatnya kemampuan invasif lokal ameloblastoma.

Kesimpulan: Ameloblastoma pleksiform dan folikuler sama-sama memiliki kemampuan invasif lokal yang sama.

.....
Background: Expression of TIMP-2 (Tissue Inhibitor of Metalloproteinase-2) immunohistochemically was used to evaluate local invasive characteristic of ameloblastomas which contributed to recurrence.

Objective: To evaluate expression of TIMP-2 in plexiform and follicular ameloblastoma.

Method: Plexiform ($n=16$) and follicular ($n=14$) ameloblastoma's samples were immunohistochemically examined with monoclonal antibody TIMP-2. Expression of TIMP-2 was evaluated with Image J software.

Result: No significant difference of immunohistochemical expression of TIMP-2 between plexiform and follicular ameloblastoma $p=0.072$ ($p>0.05$), that was analyzed with Chi-Square test.

Discussion: Low grade TIMP-2 expression was contributed to local invasive capacity of ameloblastomas.

Conclusion: Plexiform and follicular ameloblastoma have similarity in capacity of local invasiveness.