

Hubungan faktor klinis sosio demografi dan lingkungan dengan kejadian tuberkulosis paru usia 15 tahun di provinsi DKI Jakarta tahun 2013 analisis data riset kesehatan dasar 2013 = The relationship of clinical socio demography and environmental factors with the occurrence of pulmonary tuberculosis age 15 years in DKI Jakarta in 2013 data analysis of basic health research 2013

Megawati Adhitama Putri, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20412243&lokasi=lokal>

Abstrak

[ABSTRAK]

Tuberkulosis paru merupakan penyakit yang masih menjadi masalah kesehatan di dunia termasuk di Indonesia, negara dengan kasus TB terbanyak kelima di dunia. Prevalensi TB paru di Indonesia menurut Riskesdas 2013 sebesar 0,4% (400/100.000) dan salah satu provinsi yang memiliki prevalensi TB paru di atas prevalensi nasional adalah DKI Jakarta (0,6%). Kepadatan yang tinggi, merokok tembakau, diabetes, dan malnutrisi diketahui sebagai faktor-faktor yang mempengaruhi keterpaparan terhadap infeksi Mycobacterium tuberculosis dan perkembangan TB aktif.

Tujuan penelitian ini adalah untuk mengetahui hubungan faktor klinis, sosio-demografi, dan lingkungan dengan kejadian TB paru usia ≥ 15 tahun di Provinsi DKI Jakarta tahun 2013 berdasarkan data Riskesdas 2013. Penelitian ini menggunakan desain studi potong lintang. Analisis data dengan regresi logistik ganda menggunakan SPSS versi 17.0. Penderita TB paru adalah responden yang pernah didiagnosis oleh tenaga kesehatan melalui pemeriksaan dahak dan/atau foto rontgen dalam waktu ≤ 1 tahun atau memiliki semua gejala klinis TB paru.

Hasil penelitian menemukan faktor-faktor yang berhubungan dengan kejadian TB paru usia ≥ 15 tahun yaitu, diabetes melitus (POR = 3,25), status perkawinan (tidak kawin) (POR=0,44), riwayat merokok (POR=1,93), dan interaksi antara status gizi dengan umur (POR=21,31). Pengendalian dan pencegahan terhadap faktor risiko TB diantaranya edukasi tentang gizi, konseling mengenai pentingnya menjaga kadar gula darah bagi para diabetesi, dan memperluas Kawasan Tanpa Rokok penting untuk dilakukan guna menurunkan prevalensi TB paru serta mendukung keberhasilan program pengendalian TB paru di DKI Jakarta.

<hr>

<i>ABSTRACT</i>

;The purpose of this study is to determine the relationship of clinical, sociodemography, and environmental factors with the occurrence of pulmonary TB age

􀂕􀀔􀀘􀀃

􀁜􀁈􀁄􀁕􀁖􀀃

􀁌􀁑􀀃 􀀧􀀮􀀬􀀃

􀀭􀁄􀁎􀁄􀁕􀁗􀁄􀀃

􀁌􀁑􀀃 􀀕􀀓􀀔􀀖􀀃

􀁅􀁄􀁖􀁈􀁇􀀃

􀁒􀁑􀀃

􀀥􀁄􀁖􀁌􀁆􀀃
􀀫􀁈􀁄􀁏􀁗􀁋􀀃 Research 2013.

The study design is cross-sectional. Data analysis used in this study is multiple logistic regression using SPSS version 17.0. Patient with pulmonary TB is a respondent who had been diagnosed by health professionals through the examination of 􀁖􀁓􀁘􀁗􀁘􀁐􀀃 􀁄􀁑􀁇􀀒􀁒􀁕􀀃 􀁏􀁘􀁑􀁊􀀃 􀁕􀁒􀁑􀁗􀁊􀁈􀁑􀀃 􀁚􀁌􀁗􀁋􀁌􀁑􀀃 􀂔􀀔􀀃 􀁜􀁈􀁄􀁕􀀃 􀁒􀁕􀀃 􀁚􀁋􀁒􀀃 􀁋􀁄􀁇􀀃 􀁆􀁏􀁌􀁑􀁌􀁆􀁄􀁏􀀃 􀁖􀁜􀁐􀁓􀁗􀁒􀁐􀁖􀀃 􀁒􀁉􀀃 pulmonary TB.

The study result found that factors related to the occurrence of pulmonary 􀀷􀀥􀀃 􀁄􀁊􀁈􀀃 􀂕􀀔􀀘􀀃 􀁜􀁈􀁄􀁕􀁖􀀃 􀁒􀁏􀁇􀀃 􀁄􀁕􀁈􀀃 􀁇􀁌􀁄􀁅􀁈􀁗􀁈􀁖􀀃 􀀋􀀳􀀲􀀵􀀠􀀖􀀏􀀕􀀘􀀌􀀏􀀃 􀁐􀁄􀁕􀁌􀁗􀁄􀁏􀀃 􀁖􀁗􀁄􀁗􀁘􀁖􀀃 􀀋􀁘􀁑􀁐􀁄􀁕􀁕􀁌􀁈􀁇􀀌􀀃

(POR=0,44), smoking history (POR=1,93), and interaction between nutritional status and age (POR=21,31). Control and preventive efforts of risk factors of pulmonary TB, such as nutrition education, counseling about the need to control blood glucose among diabetes patient, and expanding no smoking area are important to do in order to decrease prevalence of pulmonary TB and support the success of pulmonary TB control program in DKI Jakarta, The purpose of this study is to determine the relationship of clinical, sociodemography, and environmental factors with the occurrence of pulmonary TB age 􀂕􀀔􀀘􀀃 􀁜􀁈􀁄􀁕􀁖􀀃

􀁌􀁑􀀃 􀀧􀀮􀀬􀀃
􀀭􀁄􀁎􀁄􀁕􀁗􀁄􀀃
􀁌􀁑􀀃 􀀕􀀓􀀔􀀖􀀃
􀁅􀁄􀁖􀁈􀁇􀀃
􀁒􀁑􀀃
􀀥􀁄􀁖􀁌􀁆􀀃
􀀫􀁈􀁄􀁏􀁗􀁋􀀃 Research 2013.

The
study design is cross-sectional. Data analysis used in this study is multiple logistic
regression using SPSS version 17.0. Patient with pulmonary TB is a respondent
who had been diagnosed by health professionals through the examination of
􀁖􀁓􀁘􀁗􀁘􀁐􀀃
􀁄􀁑􀁇􀀒􀁒􀁕􀀃
􀁏􀁘􀁑􀁊􀀃
􀁕􀁒􀁑􀁗􀁊􀁈􀁑􀀃
􀁚􀁌􀁗􀁋􀁌􀁑􀀃
􀂔􀀔􀀃 􀁜􀁈􀁄􀁕􀀃
􀁒􀁕􀀃 􀁚􀁋􀁒􀀃
􀁋􀁄􀁇􀀃
􀁆􀁏􀁌􀁑􀁌􀁆􀁄􀁏􀀃
􀁖􀁜􀁐􀁓􀁗􀁒􀁐􀁖􀀃 􀁒􀁉􀀃
pulmonary TB.

The study result found that factors related to the occurrence of pulmonary
􀀷􀀥􀀃 􀁄􀁊􀁈􀀃
􀂕􀀔􀀘􀀃
􀁜􀁈􀁄􀁕􀁖􀀃
􀁒􀁏􀁇􀀃 􀁄􀁕􀁈􀀃
􀁇􀁌􀁄􀁅􀁈􀁗􀁈􀁖􀀃
􀀋􀀳􀀲􀀵􀀠􀀖􀀏􀀕􀀘􀀌􀀏􀀃
􀁐􀁄􀁕􀁌􀁗􀁄􀁏􀀃
􀁖􀁗􀁄􀁗􀁘􀁖􀀃
􀀋􀁘􀁑􀁐􀁄􀁕􀁕􀁌􀁈􀁇􀀌􀀃
(POR=0,44), smoking history (POR=1,93), and interaction between nutritional
status and age (POR=21,31). Control and preventive efforts of risk factors of
pulmonary TB, such as nutrition education, counseling about the need to control
blood glucose among diabetes patient, and expanding no smoking area are
important to do in order to decrease prevalence of pulmonary TB and support the

success of pulmonary TB control program in DKI Jakarta]