

# Perbedaan fungsi seksual pada ibu postpartum pervaginam dan ibu pasca bedah sesar pada enam bulan pertama pasca melahirkan di Kabupaten Majene = Comparation sexual function of postpartum women after vaginal delivery and women caesarean section during the six months after childbirth in Majene

Immawanti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20415270&lokasi=lokal>

---

## Abstrak

[Perempuan pasca melahirkan adalah kelompok yang berisiko mengalami perubahan pada fungsi seksual. Penelitian ini untuk mengetahui perbedaan fungsi seksual ibu postpartum pervaginam dengan atau tanpa episiotomi dan pasca bedah sesar. Penelitian ini merupakan studi deskriptif dengan desain cross sectional. Sampel berjumlah 225 ibu postpartum yang diambil secara consecutive sampling dari bulan April-Mei 2015. Fungsi seksual dinilai dengan Sexual Function Questionnaire (SFQ). Hasil penelitian menunjukkan tidak ada perbedaan yang bermakna antara fungsi seksual pada semua jenis persalinan ( $p=0,977$ ), begitupun hasrat seksual, gairah seksual, dan orgasme. Penelitian ini dapat digunakan untuk meningkatkan kesehatan seksual ibu pada perawatan postpartum yang berkualitas.;The postpartum women is one of group who experience sexual function changes. The aims of the study is to know the comparation sexual function woman after vaginal delivery without episiotomi and post caesarean section. The study was a descriptive study of cross-sectional design. The sample was 225 woman

postpartum was taken by consecutive sampling from April-May 2015. Sexual function was assessed by the Sexual Function Questionnaire (SFQ) score. The results of this study showed there were no significant differences between sexual function and types of delivery ( $p=0.977$ ), including sexual desire, sexual arousal and orgasm. The results of the study can be used to increase nurses's service sexual postpartum women for reach quality postpartum care., The postpartum women is one of group who experience sexual function changes.

The aims of the study is to know the comparation sexual function woman after vaginal delivery without episiotomi and post caesarean section. The study was a descriptive study of cross-sectional design. The sample was 225 woman postpartum was taken by consecutive sampling from April-May 2015. Sexual function was assessed by the Sexual Function Questionnaire (SFQ) score. The results of this study showed there were no significant differences between sexual function and types of delivery ( $p=0.977$ ), including sexual desire, sexual arousal and orgasm. The results of the study can be used to increase nurses's service sexual postpartum women for reach quality postpartum care.]