

Isolasi dan identifikasi senyawa dari caesalpinia sappan L. lignum (kayu secang) dengan aktivitas penghambatan terhadap enzim arginase = Isolation and identification of compound from caesalpinia sappan l lignum kayu secang with arginase inhibitory activity

Arikadia Noviani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20415337&lokasi=lokal>

Abstrak

[Caesalpinia sappan L. lignum atau kayu secang adalah anggota suku Fabaceae. Kayu secang secara empiris dapat mengobati berbagai penyakit yang berkaitan dengan gangguan pembuluh darah. Penelitian sebelumnya menyatakan bahwa ekstrak kayu secang dapat menghambat aktivitas enzim arginase. Penelitian ini bertujuan untuk mengisolasi dan mengidentifikasi senyawa dengan aktivitas penghambatan terhadap enzim arginase secara in vitro. Isolasi dilakukan berdasarkan bioassay-guided isolation dengan kromatografi kolom. Dari ekstrak etil asetat didapatkan subfraksi dengan potensi penghambatan terhadap enzim arginase, yaitu subfraksi J dan K yang masing-masing mempunyai nilai IC₅₀ 67,02 μg/mL dan 75,57 μg/mL. Identifikasi golongan senyawa menunjukkan subfraksi J dan K mengandung senyawa golongan flavonoid dan terpenoid. Dari ekstrak metanol didapatkan isolat dengan IC₅₀ 265,03 μg/mL. Hasil elucidasi struktur isolat dari data spektroskopi UV-Vis, IR, MS, ¹H-NMR, ¹³C-NMR, dan 2D-NMR menunjukkan bahwa isolat tersebut adalah 2-(3,4-dihidroksifenil)-3,5,7-trihidroksikroman-4-on atau kuersetin.;Caesalpinia sappan L. lignum or kayu secang is the member of Fabaceae family. Kayu secang is empirically used as a treatment for various disease related to vascular dysfunction. Previous research reported that the extract of kayu secang was able to inhibit arginase enzyme activity. This research's aim is to isolate and identicate compound with arginase inhibitory activity in vitro. Isolation is conducted based on bioassay-guided isolation by column chromatography. From ethyl acetate extract, potential inhibition of arginase enzyme activity was exhibited by subfraction J and K with IC₅₀ 67,02 μg/mL and 75,57 μg/mL, respectively. Identification showed that subfraction J and K contains flavonoid and terpenoid compounds. From methanol extract, there is an isolate with IC₅₀ 265,03 μg/mL. Structure elucidation by spectroscopy UV-Vis, IR, MS, ¹H-NMR, ¹³C-NMR, and 2D-NMR showed that the isolate is 2-(3,4-dihidroksifenil)-3,5,7-trihidroksikroman-4-on or quercetin;Caesalpinia sappan L. lignum or kayu secang is the member of Fabaceae family. Kayu secang is empirically used as a treatment for various disease related to vascular dysfunction. Previous research reported that the extract of kayu secang was able to inhibit arginase enzyme activity. This research's aim is to isolate and identicate compound with arginase inhibitory activity in vitro. Isolation is conducted based on bioassay-guided isolation by column chromatography. From ethyl acetate extract, potential inhibition of arginase enzyme activity was exhibited by subfraction J and K with IC₅₀ 67,02 μg/mL and 75,57 μg/mL, respectively. Identification showed that subfraction J and K contains flavonoid and terpenoid compounds. From methanol extract, there is an isolate with IC₅₀

265,03 μ g/mL. Structure elucidation by spectroscopy UV-Vis, IR, MS, ¹H-NMR, ¹³C-NMR, and 2D-NMR showed that the isolate is 2-(3,4-dihydroksifenil)-3,5,7-trihidroksikroman-4-on or quercetin, *Caesalpinia sappan* L. lignum or kayu secang is the member of Fabaceae

family. Kayu secang is empirically used as a treatment for various disease related to vascular dysfunction. Previous research reported that the extract of kayu secang was able to inhibit arginase enzyme activity. This research's aim is to isolate and identify compound with arginase inhibitory activity in vitro. Isolation is conducted based on bioassay-guided isolation by column chromatography. From ethyl acetate extract, potential inhibition of arginase enzyme activity was exhibited by subfraction J and K with IC₅₀ 67,02 μ g/mL and 75,57 μ g/mL, respectively. Identification showed that subfraction J and K contains flavonoid and terpenoid compounds. From methanol extract, there is an isolate with IC₅₀ 265,03 μ g/mL. Structure elucidation by spectroscopy UV-Vis, IR, MS, ¹H-NMR, ¹³C-NMR, and 2D-NMR showed that the isolate is 2-(3,4-dihydroksifenil)-3,5,7-trihidroksikroman-4-on or quercetin]