

Intervensi keperawatan terapi hypnoparenting untuk mengatasi masalah balita dengan risiko keterlambatan perkembangan pada keluarga di Kelurahan Sukamaju Baru, Depok = Hypnoparenting as a nursing intervention to overcome toddler with risk for delayed development in family at Kelurahan Sukamaju Baru, Depok

Tissa Aulia Putri, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20416949&lokasi=lokal>

Abstrak

[Pola asuh yang tidak tepat menjadi akar masalah kesehatan balita di daerah perkotaan yang dapat mempengaruhi tumbuh kembang balita. Karya Ilmiah Akhir ini bertujuan untuk memberikan gambaran tentang asuhan keperawatan keluarga pada keluarga balita dengan masalah risiko keterlambatan perkembangan di Kelurahan Sukamaju Baru, Depok. Intervensi keperawatan unggulan adalah terapi hypnoparenting. Hasil evaluasi menunjukkan peningkatan skor kuesioner pra skrining perkembangan anak, pengetahuan, motivasi Ibu terhadap pemberian stimulasi anak dan kemandirian keluarga. Intervensi ini untuk ibu dan balita yang dilakukan selama satu bulan dengan minimal terapi 30 kali agar hasilnya optimal.;

Improper parenting became the cause of health problem in toddler in urban areas that affected their development and growth negatively. This paper aimed to depict the nursing care process within families with toddlers who had risk for delayed development in Kelurahan Sukamaju Baru, Depok. The focus of nursing care is the implementation of hypnoparenting. Result showed that there was improvements in terms of pre-screening questionnaire of development score, and both mother's knowledge and motivation towards stimulation for the toddlers, as well as family independency. This intervention for mothers and toddler carried out during for a month with a minimum of 30 times the therapy for optimal results.;

Improper parenting became the cause of health problem in toddler in urban areas that affected their development and growth negatively. This paper aimed to depict the nursing care process within families with toddlers who had risk for delayed development in Kelurahan Sukamaju Baru, Depok. The focus of nursing care is the implementation of hypnoparenting. Result showed that there was improvements in terms of pre-screening questionnaire of development score, and both mother's knowledge and motivation towards stimulation for the toddlers, as well as family independency. This intervention for mothers and toddler carried out during for a month with a minimum of 30 times the therapy for optimal results.;

Improper parenting became the cause of health problem in toddler in urban areas that affected their development and growth negatively. This paper aimed to depict the nursing care process within families with toddlers who had risk for delayed development in Kelurahan Sukamaju Baru, Depok. The focus of nursing care is the implementation of hypnoparenting. Result showed that there was improvements in terms of pre-screening questionnaire of development score, and both mother's knowledge and motivation towards stimulation for the toddlers, as well as family independency. This intervention for mothers and toddler carried out during for a month with a minimum of 30 times the therapy for optimal results.;

Improper parenting became the cause of health problem in toddler in urban areas that affected their development and growth negatively. This paper aimed to depict the nursing care process within families with toddlers who had risk for delayed development in Kelurahan Sukamaju Baru, Depok. The focus of nursing care is the implementation of hypnoparenting. Result showed that there was improvements in terms of pre-screening questionnaire of development score, and both mother's knowledge and motivation towards stimulation for the toddlers, as well as family independency. This intervention for mothers and toddler carried out during for a month with a minimum of 30 times the therapy for optimal results.;

questionnaire of development score, and both mother's knowledge and motivation towards stimulation for the toddlers, as well as family independency. This intervention for mothers and toddler carried out during for a month with a minimum of 30 times the therapy for optimal results.]