Peran S100B serum sebagai prediktor derajat keparahann trauma kepala pada anak = The role of serum S100B as predictor of severity in pediatric traumatic brain injury

Lies Dewi Nurmalia, author

Deskripsi Lengkap: https://lib.ui.ac.id/detail?id=20424627&lokasi=lokal

Abstrak

ABSTRAK

Latar Belakang Biomarker dapat digunakan untuk memprediksi derajat keparahan trauma kepala Tujuan Mengetahui hubungan antara kadar S100B dengan derajat keparahan trauma kepala dan kelainan CT scan kepala Metode Penelitian Penelitian potong lintang di IGD RSUPN Cipto Mangunkusumo RSUP Fatmawati dan RS Permata Cibubur selama Juli Desember 2015 Subjek adalah anak usia 1 18 tahun yang mengalami trauma kepala dengan onset

<hr>

ABSTRACT

Background Biomarker has ability to predict the severity of TBI and abnormal CT scan Objectives To determine the association between S100B level with the severity of pediatric TBI and intracranial injury Methods A cross sectional study at Emergency Department of RSUPN Cipto Mangunkusumo RSUP Fatmawati and Permata Cibubur Hospital on July December 2015 Subjects were 1 18 year old children with TBI onset within 24 hours before admission We measured SKG score serum S100B level and performed cranial CT scan if indicated Results Twenty subjects had mild TBI and 18 subjects had moderate TBI were included S100B levels were higher in children with moderate TBI as compared to children with mild TBI 0 173 0 054 0 812 g L vs 0 067 0 039 0 084 g L p;Background Biomarker has ability to predict the severity of TBI and abnormal CT scan Objectives To determine the association between S100B level with the severity of pediatric TBI and intracranial injury Methods A cross sectional study at Emergency Department of RSUPN Cipto Mangunkusumo RSUP Fatmawati and Permata Cibubur Hospital on July December 2015 Subjects were 1 18 year old children with TBI onset within 24 hours before admission We measured SKG score serum S100B level and performed cranial CT scan if indicated Results Twenty subjects had mild TBI and 18 subjects had moderate TBI were included S100B levels were higher in children with moderate TBI as compared to children with mild TBI 0 173 0 054 0 812 g L vs 0 067 0 039 0 084 g L p;Background Biomarker has ability to predict the severity of TBI and abnormal CT scan Objectives To determine the association between S100B level with the severity of pediatric TBI and intracranial injury Methods A cross sectional study at Emergency Department of RSUPN Cipto Mangunkusumo RSUP Fatmawati and Permata Cibubur Hospital on July December 2015 Subjects were 1 18 year old children with TBI onset within 24 hours before admission We measured SKG score serum S100B level and performed cranial CT scan if indicated Results Twenty subjects had mild TBI and 18 subjects had moderate TBI were included S100B levels were higher in children with moderate TBI as compared to children with mild TBI 0 173 0 054 0 812 g L vs 0 067 0 039 0 084 g L p;Background Biomarker has ability to predict the severity of TBI and abnormal CT scan Objectives To determine the association between S100B level with the severity of pediatric TBI and intracranial injury Methods A cross sectional study at Emergency Department of RSUPN Cipto Mangunkusumo RSUP Fatmawati and Permata Cibubur Hospital on July December 2015 Subjects

were 1 18 year old children with TBI onset within 24 hours before admission We measured SKG score serum S100B level and performed cranial CT scan if indicated Results Twenty subjects had mild TBI and 18 subjects had moderate TBI were included S100B levels were higher in children with moderate TBI as compared to children with mild TBI 0 173 0 054 0 812 g L vs 0 067 0 039 0 084 g L p