

# Proporsi kepositivan dan genotipe human papillomavirus pada suami pasien kanker serviks serta kesamaan genotipe di antara pasangan di RSUPN Dr. Cipto Mangunkusumo = The positivity proportion and genotype of human papillomavirus in cervical cancer patient's spouse and the genotype concordance between the couples at Dr. Cipto Mangunkusumo Hospital

Jonathan Raharjo Subekti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20424628&lokasi=lokal>

---

## Abstrak

**ABSTRAK**  
Latar belakang. Infeksi human papillomavirus (HPV) pada genital laki-laki selain dapat

menyebabkan kutil kelamin dan kanker penis juga meningkatkan risiko infeksi HPV pada pasangan. Walaupun saat ini telah terdapat banyak penelitian mengenai peran HPV risiko tinggi terhadap karsinogenesis serviks dan semakin jelas peran laki-laki sebagai vektor virus HPV, namun pemeriksaan HPV pada laki-laki belum rutin dilakukan. Penelitian ini bertujuan mengetahui proporsi kepositivan, variasi genotipe HPV pada suami pasien kanker serviks serta kesamaan genotipe HPV antara suami pasien kanker serviks yang HPV positif dengan pasien kanker serviks di RSUPN Dr Cipto Mangunkusumo. Metode. Penelitian potong lintang. Pemilihan SP dilakukan secara berurutan (consecutive sampling). Sampel diambil dengan menggunakan kertas amplas dan dacron swab. Pada spesimen dilakukan pemeriksaan menggunakan HPV express matrix Kalgen®. Hasil. Sebanyak 47 SP dilibatkan dalam penelitian ini, dengan rerata usia 50,7+10,6 tahun. Dari analisis spesimen diidentifikasi HPV genital pada 9 (19%) SP, terdiri atas genotipe risiko rendah (3 SP) dan risiko tinggi (6 SP). Genotipe HPV yang ditemukan adalah 6, 18, 31, 39, 43, 53, dan 56, dengan tipe tersering adalah 18 dan 43. Tidak didapatkan kesamaan tipe HPV di antara pasangan HPV yang positif. Kesimpulan. Proporsi kepositivan HPV pada suami pasien kanker serviks sebesar 19% dengan tipe 18 dan 43 paling banyak didapatkan, namun tidak didapatkan kesamaan tipe HPV antara SP dengan pasien kanker serviks pasangannya.

**ABSTRACT**  
Background. Human papillomavirus (HPV) infection on male genital could cause genital warts, penile cancer, but also increase the risk of HPV infection in their spouse. Despite many current researches on role of high-risk HPV in cervix carcinogenesis and male partner's role as HPV vector is well known, HPV examination on male is not yet routinely performed. The aim of this study is to find the positivity proportion and genotype variant of HPV on cervical cancer patient's spouse, and also the genotype concordance between the spouse with HPV positive and the cervical cancer patient at dr Cipto Mangunkusumo hospital. Method. Cross-sectional design. Subject was chosen consecutively (consecutive sampling). Sample was collected with emery paper and dacron swab. The specimen was then analyzed with HPV express matrix Kalgen®. Result. Fourty seven subject enrolled in this studi with mean age 50,7+10,6 y.o. Specimen analysis identified genital HPV on 9 (19%) subject, with low risk (3 subject) and high risk (6 subject) genotype. HPV genotypes found in this study are 6, 18, 31,

39, 43, 53, dan 56, with 18 and 43 as the most frequent. No genotype concordance found between the cervical cancer patient's spouse with HPV positive and their partners. HPV genotypes variation found on cervical cancer's spouses are type 6, 18, 31, 39, 43, 53, dan 56.

Conclusion. The positivity proportion of HPV on cervical cancer patient's spouse was 19%, with genotype 18 and 43 as the most frequent with no HPV genotype concordance found between subjects and the spouse.

Background. Human papillomavirus (HPV) infection on male genital could cause genital warts, penile cancer, but also increase the risk of HPV infection in their spouse. Despite many current researches on role of high-risk HPV in cervix carcinogenesis and male partner's role as HPV vector is well known, HPV examination on male is not yet routinely performed. The aim of this study is to find the positivity proportion and genotype variant of HPV on cervical cancer patient's spouse, and also the genotype concordance between the spouse with HPV positive and the cervical cancer patient at dr Cipto Mangunkusumo hospital.

Method. Cross-sectional design. Subject was chosen consecutively (consecutive sampling). Sample was collected with emery paper and dacron swab. The specimen was then analyzed with HPV express matrix Kalgen®.

Result. Fourty seven subject enrolled in this studi with mean age 50,7+10,6 y.o. Specimen analysis identified genital HPV on 9 (19%) subject, with low risk (3 subject) and high risk (6 subject) genotype. HPV genotypes found in this study are 6, 18, 31, 39, 43, 53, dan 56, with 18 and 43 as the most frequent. No genotype concordance found between the cervical cancer patient's spouse with HPV positive and their partners. HPV genotypes variation found on cervical cancer's spouses are type 6, 18, 31, 39, 43, 53, dan 56.

Conclusion. The positivity proportion of HPV on cervical cancer patient's spouse was 19%, with genotype 18 and 43 as the most frequent with no HPV genotype concordance found between subjects and the spouse.

Background. Human papillomavirus (HPV) infection on male genital could cause genital warts, penile cancer, but also increase the risk of HPV infection in their spouse. Despite many current researches on role of high-risk HPV in cervix carcinogenesis and male partner's role as HPV vector is well known, HPV examination on male is not yet routinely performed. The aim of this study is to find the positivity proportion and genotype variant of HPV on cervical cancer patient's spouse, and also the genotype concordance between the spouse with HPV positive and the cervical cancer patient at dr Cipto Mangunkusumo hospital.

Method. Cross-sectional design. Subject was chosen consecutively (consecutive sampling). Sample was collected with emery paper and dacron swab. The specimen was then analyzed with HPV express matrix Kalgen®.

Result. Fourty seven subject enrolled in this studi with mean age 50,7+10,6 y.o. Specimen analysis identified genital HPV on 9 (19%) subject, with low risk (3 subject) and high risk (6 subject) genotype. HPV genotypes found in this study are 6, 18, 31, 39, 43, 53, dan 56, with 18 and 43 as the most frequent. No genotype concordance found between the cervical cancer patient's spouse with HPV positive and their partners. HPV genotypes variation found on cervical cancer's spouses are type 6, 18, 31, 39, 43, 53, dan 56.

Conclusion. The positivity proportion of HPV on cervical cancer patient's spouse was 19%, with genotype 18 and 43 as the most frequent with no HPV genotype concordance found between subjects and the spouse.