

Pengaruh aromaterapi kenanga (cananga odorata) terhadap gangguan tidur insomnia pada pasien stroke di RSUD Pangkalpinang dan Sungailiat = The influence of kenanga aromatherapy (cananga odorata) to sleep disorder insomnia in patients with stroke

Eny Erlinda Widyaastuti, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20432991&lokasi=lokal>

Abstrak

ABSTRAK

Stroke dapat menyebabkan perubahan atau kerusakan neurologis berupa gangguan tidur insomnia. Gangguan tidur insomnia pada pasien pasca stroke akan mempengaruhi rehabilitasi dan kualitas hidup pasien. Oleh karena itu diperlukan suatu upaya intervensi keperawatan yang dapat meningkatkan relaksasi pasien. Penelitian ini bertujuan untuk mengetahui pengaruh aromaterapi: Kenanga (Cananga odorata) terhadap gangguan tidur insomnia pada pasien stroke di RSUD Pangkalpinang dan Sungailiat. Penelitian ini menggunakan desain kuasi eksperimen dengan pre dan posttest design dan melibatkan 38 orang responden yang dikelompokkan menjadi intervensi dan kontrol. Pemilihan responden penelitian dengan teknik consecutive sampling. Hasil Penelitian menunjukkan ada perbedaan yang signifikan rerata derajat insomnia setelah pemberian aromaterapi Kenanga antara kelompok intervensi dan kontrol ($p=0,000$). Oleh karena itu, peneliti merekomendasikan aromaterapi Kenanga sebagai salah satu alternatif intervensi keperawatan untuk masalah insomnia pada pasien stroke

ABSTRACT

Stroke may lead to altered or impaired neurological function which include insomnia. Insomnia in post-stroke patients affects the patients' rehabilitation process and quality of life. Therefore, it is of high importance to develop and examine the nursing intervention aiming at improving sleep pattern in this group of patients. The study aimed to assess the influence of Cananga (Cananga Odorata) aromatherapy to insomnia sleep disorder in patients with stroke at Pangkalpinang and Sungailiat Hospitals. This quasi-experiment with pre-posttest design was carried out in 38 respondents, consecutively sampled and assigned into the intervention and control groups. The results showed that there was a statistically significant difference between the mean of insomnia degree in the intervention and control group, after the Cananga aromatherapy treatment ($p=0.000$). It could be concluded that Cananga aromatherapy is likely to lower insomnia in patients with stroke in this study. Nurses may use this intervention to help addressing insomnia problem among patients with stroke