

Efektifitas terapi penghentian pikiran dan terapi kognitif pada klien ansietas menggunakan pendekatan model teori adaptasi Roy = Effectiveness of thought stopping and cognitive therapy to cliens anxiety approach Roy adaptation theory model

Ulfa Suryani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20435242&lokasi=lokal>

Abstrak

ABSTRAK

Penyakit fisik sering berdampak pada masalah psikologis (gangguan mental emosional). Masalah psikososial yang paling banyak terjadi adalah pada masalah depresi dan ansietas yang akan menghambat proses penyembuhan. Ansietas merupakan perasaan khawatir, tidak pasti terhadap sesuatu yang tidak jelas.

Tujuan: menjelaskan efektifitas terapi penghentian pikiran dan terapi kognitif pada klien ansietas menggunakan pendekatan model teori adaptasi Roy. Analisis dilakukan pada 26 klien penghentian pikiran+kognitif terapi, dan 24 klien penghentian pikiran. Hasil: pemberian terapi penghentian pikiran+terapi kognitif lebih meningkatkan penurunan tanda dan gejala ansietas dan meningkatkan kemampuan klien ansietas. Penggunaan pendekatan terapi penghentian pikiran mampu mengatasi pikiran irrasional yang muncul berulang sedangkan terapi kognitif dapat menggantikan pikiran otomatis negatif. Saran: kombinasi penghentian pikiran+terapi kognitif pilihan pada awal program pemulihan klien dengan penyakit fisik yang mengalami ansietas yang dapat meningkatkan hasil pengobatan.

ABSTRACT

Physical illness often affectEed the psychological problems (mental emotional disorder).Psychosocial problems was most common in depression and anxiety issues that impeded the healing process.Anxiety was a feeling worried, uncertain about something that is not clear.Objective: to explain the effectiveness of thought stopping and cognitive therapy on anxiety clients with physical illness approach Roy adaptation theory model. Analysis was conducted on 26 clients thought stopping + cognitive therapy, and 24 clients thought stopping. Results: Thought stopping and cognitive therapy further improved the signs and symptoms decreased anxiety and improved the ability of client anxiety.The used of therapeutic approaches thought stopping is able to terminated irrational thoughts that appear repeatedly while cognitive therapy can replaced negative automatic thoughts. Recomendation: a combination of thought stopping and cognitive therapy selected at the beginning of the recovery program clients with physical illness that experience anxiety that can improved treatment outcomes.;