## Periodontal status and treatment needs among the Palestinian refugees living in the united nations camps in Jordan

Deskripsi Lengkap: https://lib.ui.ac.id/detail?id=20436212&lokasi=lokal

-----

Abstrak

Indonesian Journal of Dentistry 2006; Edisi Khusus KPPIKG XIV: 25-31

A periodontal health survey, using the WHO crireria 1997. was conducted among 585 Palestinian refugees living in the United Nations Camps in Jordan, the sample consists of 363 females and 222 males aged 15-54 years. The purposes ofthh study were to determine the prevalence and the severity of periodontal disease and the treatment needs among this population and to correlate the periodontal status with age and gender. By using WHO periodontal probes, the community periodontal index (CPI), the treatment needs (TN) and the loss of attachment (LA) were measured for subject. The number of permanent teeth loss was also recorded. Females were found to have worse periodontal conditions than males. Also, a significant age difference was present. All subjects of the present population had experienced periodontal disease, where 34% and 43% of them had a shallow pocket of 4-5 mm and deep pocket of 6 mm or more respectively.

The results of this study have shown that 5.48 of sextant per subject in this population were with bleeding or higher, 4.80 with calculus or higher, 2.48 with shallow pockets or higher, and 0.88 with deep pocket. This study demonstrated that the loss of attachment increased significantly with increasing age and 42% of this population had normal level of periodontal attachment, while the rest of them (58%) had different amounts of attachment loss. Almost all subjects of this population needed oral hygiene instruction, scaling and/or removal of overhangs and 43% of them needed complex periodontal treatment. The mean number of loss permanent teeth was among this population with significant age and gender difference. Our data indicated that high prevalence and severity of periodontal disease were observed in this population require programs for both prevention and treatment.