

Studi pembentukan dna adduct 8-hidroksi-2 -deoksiganosin 8-ohdg menggunakan tert-butyl hydroquinone t-bhq secara in vitro melalui reaksi fenton like = In vitro study of dna adduct 8-hydroxy 2 - deoxyguanosine 8-ohdg formation with tert butyl hydroquinone t bhq through fenton like reaction

Annisa Nuria Rodhiani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20445316&lokasi=lokal>

Abstrak

ABSTRACT

Pada penelitian ini dilakukan studi pembentukan DNA adduct 8-Hidroksi-2 -Deoksiganosin 8-OHdG sebagai biomarker kerusakan DNA dengan mereaksikan basa DNA 2 -deoksiganosin dengan t-BHQ melalui Fenton Like Reaction Cr III dan H₂O₂ pada variasi suhu 37 C dan 60 C, pH 7,4 dan pH 8,4, dengan waktu inkubasi 7 jam dan 12 jam. Hasil adduct dianalisis menggunakan instrumen HPLC reversed phase dengan detektor UV pada panjang gelombang 254 nm. Pembentukan DNA Adduct 8-OHdG dari senyawa t-BHQ terdeteksi pada reaksi 2 -dG dan t-BHQ pada suhu 60 C waktu inkubasi 12 jam, reaksi antara 2 -dG, Cr III terdeteksi pada waktu inkubasi 7 jam, pH 8,4, 37 C, pada suhu 60 C pH 7,4 dan 8,4 serta waktu inkubasi 12 jam pada suhu 37 C dan 60 C. Reaksi antara 2 -dG, t-BHQ, H₂O₂ hanya terdeteksi pada suhu 60 C waktu inkubasi 12 jam, dan reaksi antara 2 -dG, Cr III , H₂O₂, dan 2 -dG, t-BHQ, Cr III , serta 2 -dG dengan reaksi Fenton terdeteksi baik pada waktu inkubasi 7 dan 12 jam.

<hr>

ABSTRACT

Chemistry Title In Vitro Study of DNA Adduct 8 Hydroxy 2 -Deoxyguanosine 8 OHdG Formation with Tert Butyl Hydroquinone t BHQ Through Fenton Like Reaction In this research, DNA adduct formation of 8 hydroxy 2 -Deoxyguanosine 8 OHdG as biomarkers of DNA damage which conducted by reacting the DNA bases 2 -deoxyguanosine base DNA with t BHQ through the Fenton Like Reaction Cr III and H₂O₂ with variation of temperature 37 C and 60 C , pH 7,4 and pH 8,4 , and incubation time 7 hours, and 12 hours studied. The adduct results were analyzed using instruments reversed phase HPLC with UV detector at a wavelength of 254 nm. The formation of DNA Adduct 8 OHdG of t BHQ compound is detected in the reaction of 2 -dG, t BHQ at the temperature of 60 C with incubation time at 12 hours, reaction of 2 -dG, Cr III is detected when incubation time at 7 hours, pH 8,4 and temperature at 37 C, when the temperature at 60 C with pH 7,4 and 8,4 and when incubation time is at 12 hours with the temperature at 37 C and 60 C. Reaction of 2 -dG, t BHQ, H₂O₂ only detected at the temperature of 60 C with incubation time at 12 hours, and the reaction of 2 -dG, Cr III , H₂O₂, and 2 -dG, t BHQ, Cr III , also 2 -dG with Fenton reaction are detected either when incubation time at 7 and 12 hours. Keywords Tert Butyl Hydroquinone t BHQ 8 hydroxy 2 -Deoxyguanosine 8 OHdG Fenton reaction DNA adduct HPLC.