

Penekanan bantal pasir efektif untuk klien paska kateterisasi jantung dengan komplikasi: randomized controlled trial

Sinaga, Janno, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20449118&lokasi=lokal>

Abstrak

Penelitian ini bertujuan untuk mengetahui efektifitas penekanan mekanikal bantal pasir 2,3 kg antara 2, 4, 6 jam terhadap komplikasi. Metode penelitian randomized controlled trial, dengan jumlah sampel sebanyak 90 orang. Kelompok intervensi I menggunakan bantal pasir 2,3 kg 2 jam, intervensi II 4 jam, kelompok kontrol 6 jam, pengukuran dilakukan setiap 2 jam.

Hasil penelitian tidak ada mengalami perdarahan pada semua kelompok, tidak ada perbedaan insiden haematom diantara kelompok ($p = 0,866$; $\#45; = 0,05$). Ada perbedaan rasa nyaman diantara kelompok pada observasi 4 jam ($p = 0,003$; $\#45; = 0,05$) dan observasi 6 jam ($p = 0,0005$; $\#45; = 0,05$). Rekomendasi penelitian ini adalah perlunya modifikasi Standar Prosedur Operasional penggunaan bantal pasir 2,3 kg sebagai penekan mekanikal dari 6 jam menjadi 2 jam, sebab tidak meningkatkan komplikasi, akan tetapi meningkatkan rasa nyaman klien.

<hr><i>This study was to determine the effectiveness of the mechanical suppression of sandbag 2.3 kg between the 2, 4, 6 hours against complications. The research design was randomized controlled trial study, where 90 patients as sample. A 2.3 kg sandbag was applied for two hours for the first group, four hours for the second groups, and six hours for the control groups, measurements were taken every 2 hours.

The results showed that no patient has any bleeding, not difference the incidence of hematoma between groups ($p = 0.866$; $\#45; = 0.05$). That the differences of discomfort between groups were found after 4 hours ($p = 0.003$; $\#45; = 0.05$), and after 6 hours ($p = 0.0005$; $\#45; = 0.05$). It is recommended that Standard Operational Procedure modification required from six hours into two hours in using a 2.3 kg sandbag as a mechanical pressure, because there is no increase of incidence of complications, on the otherhand an improvement of comfort level is detected.</i>