

Kompres dingin dapat menurunkan nyeri anak usia sekolah saat pemasangan infus

Puji Indriyani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20449362&lokasi=lokal>

Abstrak

Prosedur pemasangan infus dapat menimbulkan nyeri dan trauma pada anak. Tujuan penelitian ini adalah untuk mengetahui pengaruh kompres hangat dan dingin terhadap skala nyeri anak usia sekolah saat pemasangan infus. Desain penelitian yang digunakan kuasi eksperimen post test only non equivalent control group yang terbagi dalam tiga kelompok yaitu kompres hangat (15 responden), kompres dingin (15 responden), dan kontrol (15 responden). Hasil uji statistik dengan Anova menunjukkan ada perbedaan bermakna antara pengaruh pada ketiga kelompok terhadap skala nyeri ($p= 0,0001$; $= 0,05$) dan hasil uji post hock menunjukkan bahwa pemberian kompres dingin mempunyai mean difference paling besar yaitu - 4,267. Berdasarkan hasil penelitian ini maka kompres dingin lebih efektif menurunkan nyeri, oleh karena itu kompres dingin lebih disarankan untuk digunakan dalam menurunkan nyeri pada anak yang dilakukan prosedur pemasangan infus.

The Effect of Cold Compress in Reducing Pain Related to IV Cannule Insertion in Children.

Intravena canule insertion procedure could cause pain and trauma in children. The purpose of this study was to determine the effect of warm and cold compress on pain scale of school-age children IV canule insertion. The study design used a quasi-experiment post-test only non-equivalent control group, which were divided into three groups: warm compress (15 respondents), a cold compress (15 respondents) and control (15 respondents). Anova analysis showed that there were significant differences in the effect of the three groups on the pain scale ($p= 0.0001$; $= 0.05$) and post hock test results showed that administration of cold compresses has the greatest mean difference (-4.267). The result suggests that a cold compress is more effective in reduce pain, therefore, it is recommended to be applied in reducing pain related to IV canule insertion among children.