

Analisis bioekologi dan potensi pemanfaatan berkelanjutan *Graphium androcles boisduval* lepidoptera: papilionidae di Sulawesi Selatan =
Bioecological analysis and sustainable use potential of *Graphium androcles boisduval* lepidoptera papilionidae in South Sulawesi

Harlina, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20454039&lokasi=lokal>

Abstrak

ABSTRAK

Penelitian ini mengkaji aspek bioekologi dan potensi pemanfaatan berkelanjutan kupu-kupu *G. androcles* di Kawasan Taman Nasional Bantimurung- Bulusaraung TN Babul dan Taman Wisata Alam Nanggala III TWA Nanggala III , Sulawesi Selatan. Penelitian dilakukan pada bulan April 2014 ndash; Maret 2015. Untuk mengetahui ciri habitat pakan digunakan metode analisis vegetasi. Aspek biologi *G. androcles* dipelajari dengan penangkaran semi alami. Data potensi pemanfaatan *G. androcles* secara berkelanjutan diperoleh dengan metode kuesioner dan dianalisis secara deskriptif - kualitatif dalam metode analisis SWOT. Hasil Penelitian menunjukkan bahwa kelimpahan *G. androcles* tertinggi di area wisata Pattunuang 45 ekor , dan terendah di area wisata Bantimurung 16 ekor . Habitat pakan *G. androcles* didominasi oleh pohon *Canarium odoratum* INP 50,88 di Pattunuang, *Cinnamomum* sp. INP 33,8 di Bantimurung, *Ficus racemosa* INP 54,4 di Salu Tandung, dan *Ardisia purpurea* INP 50,4 di Puncak. Kupu-kupu *G. androcles* dijumpai pada bulan Juni - Nopember 2014 dengan curah hujan bulanan rendah 0 ndash; 160 mm , suhu udara 29 - 300C, kelembapan 54 - 65 , dan intensitas cahaya 33 ndash; 1180 lux. Kelimpahan *G. androcles* berkaitan dengan kondisi habitat dan ketersediaan tumbuhan pakannya. *Graphium androcles* meletakkan telur pada pucuk daun *Uvaria rufa*. Tumbuhan penghasil nektar adalah *Hibiscus rosasinensis*, *Ixora* sp., *Lantana camara*, *Dendrobium phalaenopsis*, *Clorodendrum thomsonae*, *Cromolaena odorata*, dan *Eupatorium inufolia*. Siklus hidup *G. androcles* dalam upaya penangkaran hingga mencapai tahap dewasa berkisar 46 ndash; 65 hari. Kegagalan tahap telur diakibatkan oleh serangan *Camponotus* sp. di alam, dan jamur saat di penangkaran. Serangan parasitoid *Aloeides indiscretus* dijumpai pada larva instar ketiga. Kegagalan pupa disebabkan oleh pembentukan yang tidak sempurna, dan serangan patogen sehingga pupa menghitam. Tingkat keberhasilan *G. androcles* dalam upaya penangkaran belum bisa termati hingga tahap kopulasi. Alur perdagangan *G. androcles* di kawasan TN Babul terdiri dari penangkap, pengrajin, pedagang, pengumpul, dan pembeli. Di kawasan TWA Nanggala III tidak ditemukan pengrajin. Kehadiran *G. androcles* berpotensi sebagai ajang promosi di area wisata Bantimurung. Hasil analisis SWOT menunjukkan bahwa potensi pemanfaatan *G. androcles* di TN Babul berbeda dengan di TWA Nanggala III. Di kawasan TN Babul, faktor kekuatan-peluang lebih tinggi 4,78 dibandingkan dengan kelemahan-ancaman 2,15 , sehingga pengelolaannya dapat terlaksana. Di kawasan TWA Nanggala III, faktor kelemahan - ancaman mempunyai bobot yang lebih besar 3,48 dibandingkan dengan kekuatan - peluang 1,59 , sehingga dibutuhkan strategi khusus dalam pengelolaan secara berkelanjutan. Kata kunci: Bioekologi, habitat, kupu-kupu, tumbuhan inang, penangkaran, *Graphium androcles*.

<hr />

ABSTRACT

This study examines the bioecological aspects and potential of sustainable utilization of *G. androcles* in

Bantimurung Bulusaraung National Park area Babul NP and Nanggala III Nature Park Nanggala III NP in South Sulawesi. The study was conducted on April 2014 March 2015. Vegetation analysis were performed to determine the characteristics habitat and food plants of *G. androcles*. Biological aspects of *G. androcles* were obtained through captive breeding. Data of the utilization potential of *G. androcles* were obtained by questionnaire method and analyzed in descriptive qualitative using SWOT analysis method. The results showed that the highest of *G. androcles* abundance was found at Pattunuang recreation area 45 individuals , and the lowest was at Bantimurung recreation area 16 individuals . The habitat and food plants of *G. androcles* was dominated by *Canarium odoratum* trees INP 50.88 in Pattunuang, *Cinnamomum* sp. INP 33.8 in Bantimurung, *Ficus racemosa* INP 54.4 in Salu Tandung and *Ardisia purpurea* INP 50.4 in Puncak area. *Graphium androcles* was found during at the dry season June November 2014 with low monthly rainfall 0 160 mm , air temperature 29 300C, humidity 54 65 , and light intensity 33 1180 lux. *Graphium androcles* lays its eggs on the leaves of *Uvaria rufa* Annonaceae . The nectar plants are *Hibiscus rosasinensis*, *Ixora* sp., *Lantana camara*, *Dendrobium phalaenopsis*, *Clorodendrum thomsonae*, *Cromolaena odorata*, and *Eupatorium inufolia*. The Life cycle of *G. androcles* in rearing experiment to reach adult stage ranges from 46 to 65 days. The failures of egg stages were caused by attacked by *Camponotus* sp. at nature and was fungi in captive breeding. The parasitoid attack of *Aloeides indiscretus* is found in third instar larvae. The failure of pupa is caused by imperfect formation, and pathogen attack so black pupa. The success rate of *G. androcles* at rearing experiment is still low and can not be reached until it copulates. The trading flow of *G. androcles* at Babul NP consists of catchers, craftsmen, merchants, and buyers. In the area of Nanggala III NP, no craftsmen were found. The presence of *G. androcles* has potential as a promotional and iconic event in Bantimurung recreation area. The results of SWOT analysis for the sustainable use of *G. androcles* at Babul NP differ with at Nanggala III NP. At the Babul NP area represents a higher probability strength factor 4,78 compared with threat weakness 2,15 , so that management can be accomplished. In the Nanggala III NP area, the weakness threat factor were greater weight 3,48 than the opportunity strength 1,59 , so different strategy will be needed for sustainable management. Keywords Bioecology, habitats, butterflies, hostplant, breeding experiment, *Graphium androcles*.