

Korelasi antara asupan karbohidrat, protein dan lemak dengan kadar C-reactive protein: fokus pada pasien kanker paru stadium IIIB-IV = Correlation between

Patricia Amanda, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20454812&lokasi=lokal>

Abstrak

Penelitian ini bertujuan untuk mengetahui korelasi antara asupan karbohidrat, protein dan lemak dengan kadar C-Reactive Protein pada pasien kanker paru stadium IIIB-IV. Progresivitas kanker paru dipengaruhi oleh sistem kekebalan tubuh, faktor genetik dan respon inflamasi. CRP sebagai salah satu marker inflamasi dapat diandalkan sebagai salah satu parameter untuk memprediksi perkembangan kanker. Subjek didapatkan melalui consecutive sampling yang melibatkan 49 subjek kanker paru stadium IIIB ndash; IV yang tidak sedang menjalani terapi di RS Kanker 'Dharmais'. Hasil penelitian didapatkan rerata usia 55,82 12,26 tahun, sebanyak 63,3 berjenis kelamin laki-laki. Nilai median CRP yaitu 23,82 0,30 - 207,29 mg/L. Pada penelitian ini tidak ditemukan adanya korelasi yang bermakna antara asupan karbohidrat dengan kadar CRP serum $p = 0,919$, asupan protein dengan kadar CRP serum $p = 0,257$ dan asupan lemak dengan kadar CRP serum $p = 0,986$. Kesimpulan: pada penelitian ini tidak didapatkan korelasi antara asupan karbohidrat, protein dan lemak dengan kadar CRP serum sebagai penanda perkembangan kanker.

.....The aim of the study was to determine the correlation between carbohydrate, fat and protein intake with the serum C Reactive Protein level in lung cancer patients stage IIIB ndash IV. The progression of lung cancer is influenced by immune system, genetic factors and inflammatory response, therefore CRP can be relied as one of the parameters for predicting cancer cell growth. Subjects were recruited by consecutive sampling, 49 subjects with lung cancer stage IIIB IV who currently not receiving any treatment in Dharmais Cancer Hospital participating in this study. The mean age of subject was 55,82 12,26 years old and 63,3 were male. The median value of CRP was 23,82 0,30 207,29 mg L. This study did not showed significant correlation between carbohydate, protein and fat intake with serum CRP value $p 0,919$ $p 0,257$ $p 0,986$, respectively. In conclusion, there is no correlation between carbohydrate, protein and fat intake with serum CRP level in lung cancer stage IIIB ndash IV.