

## Perbedaan kebutuhan perawatan suportif pasien kanker ginekologi di Rawat Inap dan Rawat Jalan = The difference of supportive care needs in inpatient and outpatient gynecologic cancer patient

Tina Mawardika, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20454815&lokasi=lokal>

---

### Abstrak

<b>ABSTRAK</b><br>

Nama : Tina Mawardika Program Studi : Magister Ilmu Keperawatan Universitas Indonesia Judul : Perbedaan Kebutuhan Perawatan Suportif Pasien Kanker Ginekologi di Rawat Inap dan Rawat Jalan. Pelayanan kesehatan yang diterima di rawat inap dan rawat jalan dapat memunculkan perbedaan kebutuhan perawatan suportif. Penelitian cross-sectional ini bertujuan untuk mengidentifikasi perbedaan antara kebutuhan perawatan suportif pada pasien kanker ginekologi di rawat inap dan rawat jalan. Teknik consecutive sampling dilakukan untuk memilih 200 pasien kanker ginekologi. Hasil penelitian menunjukkan bahwa sebagian besar 83 pasien di rawat inap memerlukan kebutuhan perawatan suportif terutama pada domain fisik 80 dan domain psikologi 84 dan 40 pasien di rawat jalan memerlukan kebutuhan perawatan terutama pada domain sistem informasi kesehatan 78. Terdapat perbedaan yang signifikan secara statistik antara kebutuhan perawatan pada pasien kanker ginekologi di rawat inap dan rawat jalan ? value = 0,000 p le; 0,05 dan nilai OR 43,9, maka dapat disimpulkan pasien di rawat inap kebutuhan perawatan suportifnya meningkat 43,9 kali dibandingkan dengan pasien di rawat jalan. Penilaian kebutuhan perawatan suportif berguna untuk mengevaluasi kebutuhan perawatan suportif pasien kanker ginekologi di rawat inap maupun rawat jalan sehingga intervensi yang diberikan tepat guna dan tepat sasaran. Kata Kunci: Kebutuhan perawatan suportif, status perawatan, kanker ginekologi.

<hr />

<b>ABSTRACT</b><br>

Name Tina Mawardika Program Study Faculty Of Nursing Indonesia University Title The difference of Supportive Care Needs in inpatient and outpatient gynecologic cancer patient. Health services that performed at inpatient and outpatient care can cause different supportive care needs. This cross sectional study aims to identify the differences of supportive care needs in inpatient and outpatient gynecologic cancer patient. Consecutive sampling techniques were undertaken to select 200 gynecological cancer patients. The results showed that of patients undergoing hospitalization required supportive care needs, especially in the physical 80 and pshycologic domain 84, 40 of patients who underwent outpatient care require treatment especially in the information system domain 78. According to statistical result, there was a different between inpatients and outpatients supportive care needs value 0,000 p le 0.05 OR 43,9. It means that inpatient rsquo s supportive care needs increase 43,9 compared with outpatient supportive care needs. Assessment of care needs is useful to evaluate the supportive care needs of gynecological cancer patients not only inpatient but also outpatient. That rsquo s way the intervention is given efficient and on target. Key Words Supportive Care Needs, Care Status, Gynecology Cancer.