

## Variasi spasial temporal suhu permukaan daratan kota metropolitan Bandung raya tahun 2014-2016 = Spatio temporal variation of land surface temperature of Metropolitan Bandung Raya 2014-2016

Safirah Timami, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20457782&lokasi=lokal>

---

### Abstrak

Kota Metropolitan Bandung Raya mengalami perkembangan cukup pesat yang menyebabkan berkurangnya lahan bervegetasi. Selain itu, perbedaan curah hujan pada setiap musim di tahun 2014 ndash; 2016 dapat mempengaruhi tingkat kehijauan vegetasi. Minimnya lahan bervegetasi menunjukkan tingkat kehijauan yang rendah sehingga dapat memicu kenaikan suhu permukaan daratan. Pengolahan citra Landsat 8/OLI dengan menggunakan algoritma Land Surface Temperature LST untuk mengetahui suhu permukaan daratan dan Nomalized Difference Index Vegetation NDVI untuk kehijauan vegetasi. Data curah hujan harian selama 15 hari sebelum waktu perekaman citra dikaji untuk menunjukkan variasi kehijauan vegetasi dan suhu permukaan daratan. Analisis dilakukan dengan menggunakan metode overlay peta dan perhitungan statistik. Hasil penelitian menunjukkan variasi suhu permukaan daratan di bagian utara merupakan wilayah suhu rendah. Pada bagian tengah merupakan wilayah dengan suhu tinggi, sementara di bagian selatan merupakan wilayah suhu yang lebih rendah. Suhu permukaan daratan yang diperoleh paling rendah ialah 0,5 dan paling tinggi >35 C dengan nilai NDVI.

*Metropolitan Bandung Raya experienced a rapid growth that caused the decreasing of vegetated land. In addition, rainfall differences in each season in 2014 2016 can affect the greenness of vegetation. The lack of vegetated land shows a low greenishness that can trigger an increase in surface temperatures. Landsat 8 OLI image processing using Land Surface Temperature LST algorithm to determine surface temperature of the land and Nomalized Difference Index Vegetation NDVI for green vegetation. Daily rainfall data for 15 days before the image recording time is examined to show variations of green vegetation and surface temperatures. The analysis is done by using map overlay and statistical calculation. The results showed that the temperature of the mainland surface in the north is the low temperature region. In the middle is a region with high temperatures, while in the southern part is the region of lower temperatures. The lowest surface temperature of the land obtained is 0,5 and at most 35 C with NDVI value.*