

Hubungan antara kematian dengan durasi kelebihan cairan pada pasien yang mendapat terapi cairan di Ruang Rawat Intensif Rumah Sakit Ciptomangunkusumo = Association between mortality and fluid overload duration in Dr. Cipto Mangunkusumo National General Hospital's Intensive Care Unit Patients

Eka Pramudita, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20465373&lokasi=lokal>

Abstrak

ABSTRAK

Latar Belakang: Resusitasi cairan merupakan terapi yang sering diberikan pada ruang rawat intensif untuk mengembalikan perfusi jaringan. Namun, seringkali terapi resusitasi cairan menyebabkan kelebihan cairan yang memiliki efek buruk terhadap pasien termasuk kematian Tujuan: Penelitian retrospektif ini bertujuan untuk mengetahui hubungan antara mortalitas dengan durasi kelebihan cairan di Rumah Sakit

Ciptomangunkusumo Metode: Sebanyak 194 pasien yang mengalami kelebihan cairan dan berada di ruang rawat intensif selama 7 hari atau lebih, diperoleh melalui teknik consecutive sampling, dievaluasi. Durasi kelebihan cairan dan kematian 28 hari dicatat. Sampel yang diperoleh dibagi menjadi dua kelompok, yaitu pasien yang mengalami kelebihan cairan kurang dari sama dengan 4 hari dan pasien yang mengalami kelebihan cairan lebih dari 4 hari. Sampel kemudian dianalisis menggunakan uji bivariat Chi square untuk diketahui hubungannya dengan kematian. Hasil: Terdapat hubungan antara kematian dengan durasi kelebihan cairan dengan nilai P.

<hr>

ABSTRACT

Background Fluid resuscitation is a common therapy given at the Intensive Care Unit ICU to maintain tissue perfusions. However, this therapy usually results in fluid overload that has adverse outcome including death Objective This retrospective study aimed to assess the association between mortality and fluid overload duration in Dr. Cipto Mangunkusumo National General Hospital 39 s Methods A total of 194 ICU patients with fluid overload and stayed for 7 days or more that obtained by consecutive sampling, were evaluated. Fluid overload duration and 28 days mortality were recorded. Samples were divided into two groups, patients with fluid overload less than or equal to 4 days and patients with fluid overload more than 4 days. A bivariate analysis Chi square were perform to assess the association of mortality and fluid overload duration. Results Mortality and fluid overload duration were significantly associated P