

Aktivitas anti-inflamasi jamu anti-aterosklerosis dengan metode in vivo pada tikus putih jantan *rattus norvegicus* = Anti inflammatory activity test of anti atherosclerosis herbs using in vivo method in white male rats *rattus norvegicus*

Riska Amalia, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20472309&lokasi=lokal>

Abstrak

ABSTRAK

Aterosklerosis merupakan kondisi pembentukan plak atau ateroma di dalam arteri yang mengandung darah kaya oksigen sehingga mempersempit dan menyebabkan kejang pada arteri. Pembentukan plak berkelanjutan mampu mengarah pada inflamasi akut, inflamasi kronis, dan aterosklerosis. Kombinasi herbal yang terdiri atas daun tanjung *Mimusops elengi* L., daun belimbing manis *Averrhoa carambola* L., dan temulawak *Curcuma xanthorrhiza* R. dipercaya secara empirik berperan sebagai jamu anti-aterosklerosis yang mengandung quercetin sebagai flavonoid dengan potensi anti-inflamasi. Pengujian aktivitas anti-inflamasi dilakukan secara in vivo pada tikus putih jantan galur Sprague-Dawley yang diinduksi oleh karagenan 1 dan terbagi dalam 5 kelompok pengujian, yaitu kontrol negatif diinduksi dan diberikan Na CMC 0,5, kontrol positif diinduksi dan diberikan Natrium diklofenak 2, dosis I diinduksi dan diberikan ekstrak jamu 2,7 mL/200 g BB, dosis II diinduksi dan diberikan ekstrak jamu 3,6 mL/200 g BB, dan dosis III diinduksi dan diberikan ekstrak jamu 4,5 mL/200 g BB. Aktivitas anti-inflamasi dari hasil ekstraksi 24 gram jamu berukuran serbuk 60.

<hr>

ABSTRACT

Atherosclerosis is plaque or atheroma formation inside rich oxygen artery which leads to narrowing and shocking the artery itself. Plaque continuous formation may lead into acute and chronic inflammation that will responsible for atherosclerosis formation. Herbs combination consists of tanjung leaves *Mimusops elengi* L., starfruit leaves *Averrhoa carambola* L., and curcuma *Curcuma xanthorrhiza* R. are empirically believed to contribute as anti atherosclerosis herbs which contains quercetin as potential flavonoid with anti inflammatory activity. Anti inflammatory activity test is done using in vivo method in Sprague Dawley strains of white male rats that carrageenan 1 induced and divided in 5 groups, which are negative control induced and given Na CMC 0.5, positive control induced and given Sodium diclofenac 2, dose I induced and given 2.7 mL 200 g BB of herbs extract, dose II induced and given 3.6 mL 200 g BB of herbs extract, and dose III induced and given 4.5 mL 200 g BB of herbs extract. 24 gram of herbs in 60.