

Hubungan tingkat depresi dengan kualitas tidur pasien gagal ginjal terminal yang menjalani hemodialisis = The relationship between depression and sleep quality in patients on end stage renal disease with hemodialysis

Nadya Rachmayani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20473829&lokasi=lokal>

Abstrak

Kualitas tidur yang baik merupakan salah satu faktor kualitas hidup yang baik. Stres emosional seperti depresi menjadi penyebab kualitas tidur buruk. Depresi dan kualitas tidur secara tidak langsung dapat mempengaruhi kejadian morbiditas dan mortalitas penyakit gagal ginjal terminal. Tujuan penelitian ini adalah mengidentifikasi hubungan antara tingkat depresi dengan kualitas tidur pasien gagal ginjal terminal yang menjalani hemodialisis. Penelitian ini merupakan penelitian deskriptif korelasi dengan metode cross sectional. Sampel yang digunakan sebanyak 92 responden dengan teknik purposive sampling. Dengan analisis bivariat menggunakan uji spearman, korelasi antara tingkat depresi dengan kualitas tidur tidak bermakna dengan nilai p 0,332. Penelitian ini menyimpulkan bahwa mayoritas responden tidak memiliki tanda klinis depresi dan kualitas tidur baik. sehingga perlu dikembangkan intervensi yang dapat meningkatkan kualitas tidurnya.

.....

Good sleep quality is one of the factors of quality of life. Emotional stress such as depression causes poor sleep quality. Depression and sleep quality indirectly may affect the incidence of morbidity and mortality of end stage renal disease. The purpose of this study was to identify the relation between depression and sleep quality in patients on end stage renal disease with hemodialysis. This research is was descriptive and correlation design with cross sectional study. There were 92 respondents that were selected by purposive sampling method. The study result shows correlation between depression and sleep quality not significant with p value 0,332 that were analysed by bivariate analysis and spearman test. The study conclude that the majority of respondents had no clinical signs of depression and good sleep quality. Therefore, it is recommended to develop nursing interventions that can improve sleep quality.