

Pengaruh edukasi manajemen nutrisi kepada orang tua terhadap mual-muntah dan perubahan berat badan pada anak dengan leukemia = The effect of nutritional management education on parents to the nausea-vomiting and weight changes in children with leukemia

Fiorentina Nova Fiorentina Nova, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20476172&lokasi=lokal>

Abstrak

ABSTRAK

Leukemia merupakan kanker yang terjadi pada pembentukan sel darah. Manajemen terapi pada anak dengan leukemia adalah kemoterapi. Chemotherapy Induced Nausea and Vomiting CINV merupakan masalah yang sering terjadi pada anak dengan pengobatan kemoterapi dan dapat mempengaruhi berat badan anak. Metode yang digunakan pada penelitian kuantitatif ini adalah quasi experiment dengan pendekatan pretest and posttest nonequivalent control group. Responden dalam penelitian ini diambil dengan cara consecutive sampling dengan jumlah sampel minimal 28 anak dengan leukemia yang sedang menjalani kemoterapi, rentang usia anak 3-18 tahun yang terbagi menjadi kelompok perlakuan n= 14 dan kelompok kontrol n=14 . Data yang diambil pada penelitian ini adalah skor mual-muntah anak INVR dan perubahan berat badan anak yang diperoleh melalui pengukuran sebelum dan sesudah intervensi edukasi dengan rentang pengukuran 7 hari. Pengetahuan orang tua pada kelompok perlakuan dilakukan pengukuran sebelum dan sesudah intervensi edukasi. Intervensi edukasi manajemen nutrisi diberikan kepada orang tua dari anak dengan leukemia melalui video yang berdurasi 10-15 menit sebelum kemoterapi diberikan. Hasil analisis menunjukkan bahwa edukasi manajemen nutrisi yang diberikan kepada orang tua tidak memiliki pengaruh yang signifikan terhadap skor mual-muntah antara sebelum dan setelah intervensi p value>0,05 . Dan pada hasil analisis skor mual-muntah setelah intervensi dan perubahan berat badan antara kelompok perlakuan dan kelompok kontrol tidak menunjukkan adanya perbedaan yang signifikan pula p value>0,05 . Hasil yang tidak signifikan pada penelitian ini terjadi disebabkan oleh faktor agen kemoterapi yang memiliki hubungan yang bermakna terhadap skor mual-muntah setelah intervensi dan variabel perubahan berat badan. Melalui hasil penelitian diketahui bahwa terjadi peningkatan pengetahuan orang tua yang mendapatkan intervensi edukasi manajemen nutrisi.

ABSTRACT

Leukemia is a cancer that occurs in the production of blood cells. Therapeutic management in the children with leukemia is chemotherapy. Chemotherapy Induced Nausea and Vomiting CINV is common problem in children with chemotherapy and may affect to the children's weight. The method was used in this quantitative study is quasi experiment with pretest and posttest nonequivalent control group. The respondents in this study were taken by consecutive sampling and the minimum sample size is 28 children with leukemia who are undergoing chemotherapy treatment with an age range of 3 to 18 years old that divided into treatment group n 14 and control group n 14 . The data that was collected in this study were nausea and vomiting scores as measured by Instrument of Nausea and Vomiting Retching INVR and children's weight changes that was measured before the education was given and 7 days later. Parent's knowledge of nutritional management in the treatment group was measured before and after

intervention. The interventions about nutritional management are given to parents of the children with leukemia through the video on the duration of 10 until 15 minutes before chemotherapy was administered. The results of the analysis showed that nutritional management education that was given to parents did not have a significant effect on the scoring of nausea vomiting between before and after intervention p value 0,05 . And on the result of the analysis to nausea vomiting scoring after intervention and weight changes between the treatment and control group is also did not show any significant differences p value 0,05 . Insignificant results on this study occurred due to chemotherapy agent factors that had a significant relationship to the score of nausea vomiting after intervention and weight changes variable. Through the result of the study is known that there are increasing in parent rsquo s knowledge who got nutritional management education intervention.