

## Determinan stunting di Indonesia Timur = The determinant of stunting in East Indonesia

Arnil Dedy Djara, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20477545&lokasi=lokal>

---

### Abstrak

#### <b>ABSTRAK</b><br>

Usia dini 0-6 tahun adalah salah satu periode paling penting bagi pertumbuhan dan perkembangan anak. Periode ini penting untuk menempatkan perkembangan kesehatan dan intelektual untuk masa depan seorang anak. Berdasarkan itu, masalah gizi pada masa anak-anak perlu dihilangkan. Menurut Riset Kesehatan Dasar Kementerian Kesehatan RI 2013, terdapat 37.2 anak yang dikategorikan kerdil atau stunting. Indonesia merupakan negara dengan stunting terbesar di ASEAN. Tujuan dari penelitian ini adalah: 1 Mengidentifikasi faktor-faktor yang mempengaruhi stunting pada anak usia dini 0-6 tahun di Indonesia timur; 2 Mengidentifikasi seberapa besar pengaruh faktor-faktor tersebut terhadap stunting pada anak usia dini 0-6 tahun di Indonesia timur. Penelitian ini menggunakan data sekunder yang diambil dari Indonesia Family Life Survey East IFLS EAST 2012. Desain yang digunakan dalam penelitian ini adalah desain potong lintang Cross Section. Sampel yang digunakan adalah anak usia 0-6 tahun berjumlah 1.515 orang. Analisis determinan stunting dilakukan menggunakan analisis regresi logistik dengan program Stata 12.0. Hasil penelitian menunjukkan bahwa faktor-faktor yang mempengaruhi stunting secara statistik adalah usia, wilayah tempat tinggal dan sumber air minum.

<hr>

#### <b>ABSTRACT</b><br>

Early age 0 6 years is one of the most important periods for the growth and development of children. This period is important to put the health and intellectual development for the future of a child. Based on that, nutritional problems in childhood need to be eliminated. According to Paimary Health Research of Health Ministry RI 2013, there are 37.2 of children who are categorized as stunting. Indonesia is the largest stunting country in ASEAN. The objectives of this study are 1 to identify the determinant factors of stunting in early childhood 0 6 years 2 to identify the probability effect of relationship between the factors and stunting in early childhood 0 6 years. This study uses secondary data from Indonesia Family Life Survey East IFLS EAST 2012 with cross sectional design. The sample size is 1.515 children aged 0 6 years. To identify the determinant factors of stunting, it uses multiple logistic regression analysis using Stata 12.0 program. The result shows that factors that effect stunting statistically are age, living place and drinking water resources.