

Efek pemberian sitikolin pada etambutol optik neuropati: analisis histopatologi & imunohistokimia tingkat kerusakan sel ganglion retina pada tikus model = Citicoline effects on ethambutol optic neuropathy: histopathological & immunohistochemical analysis of the level of retinal ganglion cell damage in rat model

Rani Indira Sari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20485463&lokasi=lokal>

Abstrak

Tujuan: Menganalisis efek pemberian sitikolin terhadap kerusakan sel ganglion tikus pada Etambutol Optik Neuropati (EON).

Metode: Penelitian ini merupakan penelitian eksperimental dengan hewan coba tikus. Lima belas ekor tikus Wistar dibagi dalam tiga kelompok yaitu kelompok tanpa perlakuan (A), kelompok etambutol (B), kelompok etambutol dan sitikolin (C). Kelompok B dan C diberikan perlakuan selama 30 hari, kemudian dilakukan pemeriksaan histopatologi untuk menilai densitas sel ganglion retina, dan imunohistokimia untuk menilai ekspresi bcl-2 dan caspase-3.

Hasil: Densitas sel ganglion retina tikus dengan intoksikasi etambutol yang mendapat sitikolin lebih tinggi dibandingkan yang tidak mendapatkan sitikolin ($p=0,001$). Lapisan ganglion tikus yang mendapat sitikolin lebih tipis dibandingkan yang tidak mendapatkan sitikolin ($p=0,005$), peningkatan ketebalan lapisan sel ganglion ini karena pembentukan vakuola pada sitoplasma sel ganglion. Tikus dengan sitikolin didapatkan ekspresi bcl-2 ganglion yang lebih tinggi ($p=0,001$), dan ekspresi caspase-3 yang lebih rendah ($p=0,02$) dibandingkan tikus yang tidak mendapatkan sitikolin.

Simpulan: Sitikolin memberikan efek proteksi terhadap sel ganglion retina dengan EON, dinilai dari morfologi sel ganglion dan ekspresi caspase-3 dan bcl-2.

<hr><i>Objective: To analyze the effects of citicoline administration on rat ganglion cell damage in Ethambutol Optic Neuropathy (EON).

Method: This study was an experimental study with rat experiments. Fifteen Wistar rats were divided into three groups, the non-treatment group (A), the ethambutol group (B), the ethambutol and citicoline group (C). Group B and C were given treatment for 30 days, then histopathological examination was performed to assess retinal ganglion cell density, and immunohistochemistry to assess bcl-2 and caspase-3 expression.

Result: retinal ganglion cell density of rat with ethambutol intoxication that received citicoline were higher than those who did not get citicoline ($p = 0.001$). The rat ganglion layer that received citicoline was thinner than those who did not get citicoline ($p = 0.005$), the increase in thickness of the ganglion cell layer was due to the formation of vacuoles in the cytoplasm of ganglion cells. Rat with citicoline obtained higher bcl-2 ganglion expression ($p = 0.001$), and lower caspase-3 expression ($p = 0.02$) than rat that did not get citicoline.

Conclusions: Citicoline have a protective effect on retinal ganglion cells with EON, judged by the morphology of ganglion cells and caspase-3 and bcl-2 expressions.</i>