

Perbandingan volume lambung setelah puasa 6 Jam dan puasa 8 Jam pada pasien yang akan menjalani pembedahan elektif = Comparison between gastric volume 6-hour and 8-hour after solid intake in patient scheduled for elective surgery

Meliani Anggreni, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20490470&lokasi=lokal>

Abstrak

Latar Belakang. Puasa sebelum pembiusan merupakan hal yang lazim dilakukan untuk menghindari aspirasi perioperatif. Lama puasa dapat menentukan volume lambung yang tersisa. Puasa yang terlalu singkat dapat meningkatkan risiko aspirasi, namun puasa yang berkepanjangan dapat menyebabkan dehidrasi dan hipoglikemia. Penelitian ini bertujuan untuk menganalisa volume lambung setelah puasa 6 jam dan 8 jam setelah makan padat pada pasien yang akan menjalani operasi elektif. Metode. Penelitian ini merupakan studi kohort terhadap 37 subjek penelitian selama Januari hingga Februari 2019. Subjek penelitian adalah pasien yang akan menjalani pembedahan elektif non-disgestif di RSUPN Dr. Cipto Mangunkusumo dengan usia antara 18-60 tahun, tidak mengalami kelainan status gizi, dinilai dengan status fisik ASA 1 atau 2. Kriteria penolakan adalah pasien dengan penyakit diabetes mellitus, kehamilan, distensi abdomen, riwayat dispepsia dan gangguan motilitas usus. Hasil. Pada penelitian ini, didapatkan volume lambung setelah puasa makan padat 6 jam adalah $35,07 \pm 39,17$ dan setelah puasa 8 jam adalah $14,16 \pm 19,24$. Selisih antara kedua rerata tersebut adalah $20,91 \pm 38,60$, $p=0,002$. Setelah puasa 6 jam, 5,4% subjek memiliki volume lambung di atas 1,5 ml/kg, sedangkan setelah puasa 8 jam, volume lambung seluruh pasien di bawah 1,5 ml/kg. Simpulan. Volume lambung setelah puasa 6 jam lebih kecil secara signifikan dibandingkan dengan puasa 8 jam.

.....Background. Preoperative fasting was a common practice to decrease perioperative aspiration risk. Duration of fasting was proportional to gastric volume. Short fasting duration may increase aspiration risk. However, prolonged perioperative fasting duration may lead to dehydration and hypoglycemia. The objective of this study was to analyze gastric volume after 6-hour and 8-hour duration of fasting after consumption of solid meal in patient scheduled for elective surgery. Methods. This was a cohort study for 37 subjects from January to February 2019. Subjects were patient scheduled for elective non-digestive surgery in RSUPN Dr. Cipto Mangunkusumo, with age between 18 to 60 years old, no nutritional status disorder and physical status of ASA 1 or 2. Exclusion criteria were patients with diabetes mellitus, pregnancy, abdominal distention, history of dyspepsia and intestinal motility disturbances. Results. In this study, gastric volume 6-hour after solid intake was 35.07 ± 39.17 and gastric volume 8-hour after solid intake was 14.16 ± 19.24 . Difference of gastric volume between 6-hour and 8-hour after solid intake was 20.91 ± 38.60 , $p=0.002$. After 6-hour of fasting, 5.4% of the subjects had gastric volume above 1,5ml/kg, while after 8-hour of fasting, gastric volume of all subjects were below 1,5ml/kg. Conclusion. Gastric volume 8-hour after solid intake was smaller than gastric volume 6-hour after solid intake.