

Hubungan tingkat pengetahuan manajemen diet dengan rasio lingk pinggang pinggul pada penderita diabetes melitus tipe 2 = Relationship between knowledge management level of diet and waist to hip ratio in people with type 2 diabetes mellitus

Zaenudin, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20492995&lokasi=lokal>

Abstrak

Sebagian besar penderita diabetes melitus tipe 2 tergolong obesitas berdasarkan pengukuran waist to hip ratio (RLPP). Obesitas sentral dapat dipengaruhi oleh kurangnya pengetahuan penderita diabetes melitus tipe 2 tentang pengelolaan diet diabetes. Penelitian ini bertujuan untuk mengetahui hubungan antara tingkat pengetahuan tentang manajemen pola makan dengan rasio lingk pinggang pinggul pada pasien diabetes melitus tipe 2. Penelitian ini menggunakan pendekatan cross-sectional dengan melibatkan 141 penderita diabetes tipe 2 yang dipilih dengan teknik consecutive sampling. Hasil penelitian ini menunjukkan bahwa sebagian besar penderita DM tipe 2 memiliki tingkat pengetahuan sedang (59,6%) dan mengalami obesitas berdasarkan pengukuran RLPP (70,2%). Hasil analisis bivariat menunjukkan bahwa ada hubungan yang signifikan antara tingkat pengetahuan tentang pengelolaan pola makan dengan rasio lingk pinggang pinggul pada penderita diabetes melitus tipe 2 (p value: 0,000). Peneliti merekomendasikan peningkatan tingkat pengetahuan tentang manajemen pola makan pada penderita diabetes melitus tipe 2, terutama tentang jenis makanan yang perlu dikonsumsi sehari-hari. Selain itu, perlu dilakukan pengukuran rasio lingk pinggang pada penderita diabetes melitus tipe 2 sehingga nilai normal rasio lingk pinggang-pinggul adalah <0,90 untuk pria dan <0,85 untuk wanita.

.....Most people with type 2 diabetes mellitus are classified as obese based on the measurement of waist to hip ratio (RLPP). Central obesity can be influenced by the lack of knowledge of type 2 diabetes mellitus sufferers about diabetes diet management. This study aims to determine the relationship between the level of knowledge about diet management and the waist-to-hip ratio in patients with type 2 diabetes mellitus. This study used a cross-sectional approach involving 141 type 2 diabetes patients who were selected by consecutive sampling technique. The results of this study indicate that most of the type 2 diabetes mellitus sufferers have a moderate level of knowledge (59.6%) and are obese based on the RLPP measurement (70.2%). The results of the bivariate analysis showed that there was a significant relationship between the level of knowledge about diet management and the waist-to-hip circumference ratio in type 2 diabetes mellitus sufferers (p value: 0.000). Researchers recommend increasing the level of knowledge about diet management in people with type 2 diabetes mellitus, especially about the types of food that need to be consumed daily. In addition, it is necessary to measure the waist circumference ratio in type 2 diabetes mellitus sufferers so that the normal value of the waist-to-hip circumference ratio is <0.90 for men and <0.85 for women.