

# Efektivitas Latihan Penguatan Faring, Latihan Pergerakan Hiolaring dan Praktik Menelan Terhadap Fungsi Menelan Penderita Stroke Iskemik dengan Disfagia Neurogenik = The Effectivity of Pharyngeal Strengthening Exercise, Hyolaryngeal Complex Range of Motion Exercise and Swallowing Practice in Swallowing Function of Ischemic Stroke Patients with Neurogenic Dysphagia

Nur Indah Lestari, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20493741&lokasi=lokal>

---

## Abstrak

Disfagia sangat berhubungan dengan peningkatan risiko pneumonia aspirasi yang sering mengakibatkan kematian pada stroke. Oleh karena itu, manajemen yang efektif dan efisien menjadi penting. Terapi perilaku rehabilitasi menelan yang berdasarkan prinsip neuroplastisitas seperti latihan penguatan dan latihan pergerakan orofaring menjadi alternatif yang cukup sering digunakan. Penelitian ini bertujuan untuk menilai perubahan fungsi menelan pada penderita stroke iskemik dengan disfagia neurogenik setelah dilakukan latihan penguatan faring, latihan pergerakan hiofarining dan praktik menelan. Fungsi menelan dinilai dengan menggunakan Penetration Aspiration Scale (PAS) dan Functional Oral Intake Scale (FOIS) berdasarkan pemeriksaan Flexible Endoscopic Evaluation of Swallowing (FEES) sebelum dan setelah intervensi. Intervensi diberikan setiap hari dengan durasi 30-45 menit selama 4 minggu. Terdapat 6 subjek yang menyelesaikan penelitian. Nilai PAS sebelum intervensi adalah  $6 \pm 1.79$  dan setelah intervensi adalah  $1.67 \pm 0.82$  ( $p=0.003$ ). Sementara itu, nilai FOIS sebelum intervensi adalah 3 (1-5) dan setelah intervensi adalah  $5 \pm 2.10$  ( $p=0.041$ ). Terdapat perbaikan nilai PAS dan FOIS setelah intervensi. Oleh karena itu, intervensi ini bisa disarankan sebagai salah satu tatalaksana dalam meningkatkan fungsi menelan pada penderita stroke iskemik dengan disfagia neurogenik.

.....Dysphagia is associated with an increased risk of aspiration pneumonia which often results in death in stroke patients. Therefore, effective and efficient management is important. Behavioral therapy for swallowing rehabilitation based on the principles of neuroplasticity such as oropharyngeal strengthening and range of motion exercises are the alternative ones that often be used. This study aimed to assess the changes in swallowing function in ischemic stroke patients with neurogenic dysphagia after pharyngeal strengthening exercise, hyolaryngeal complex range of motion exercise and swallowing practice. Swallowing function was assessed using Penetration Aspiration Scale (PAS) and Functional oral Intake Scale (FOIS) based on Flexible Endoscopic Evaluation of Swallowing (FEES) before and after interventions. The interventions were given every day with a duration of 30-45 minutes for 4 weeks. There were 6 subjects who completed the study. The PAS before the interventions was  $6 \pm 1.79$  and after the interventions was  $1.67 \pm 0.82$  ( $p=0.003$ ). Meanwhile, the FOIS score before the interventions was 3 (1-5) and after the interventions was  $5 \pm 2.10$  ( $p=0.041$ ). There was an improvement of PAS and FOIS after the interventions. Therefore, the interventions can be suggested to be used as one of the treatments to improve swallowing function in ischemic stroke patients with neurogenic dysphagia.