

Ekspresi Programmed Death Ligand-1 (PD-L1) pada Seminoma Testis sebagai Potensi Faktor Prognostik = The Expression of Programmed Death Ligand-1 (PD-L1) in Testicular Seminoma as Potential Prognostic Factor.

Freciyana, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20494079&lokasi=lokal>

Abstrak

ABSTRAK

Latar belakang: Seminoma testis yang gagal sembuh dengan penatalaksanaan konvensional memiliki prognosis buruk. Beberapa kejadian rekuren setelah dilakukan kemoterapi juga ditemukan. Programmed Death Ligand-1 (PD-L1) terekspresi pada berbagai keganasan dan tumor infiltrating lymphocytes(TILs) serta telah diketahui perannya sebagai faktor prognostik. Penelitian ini dilakukan untuk mengetahui peran ekspresi PD-L1 pada seminoma testis dalam menentukan overall survival(OS) dan progression free survival(PFS).

Bahan dan cara: Penelitian ini merupakan penelitian kohort retrospektif dengan desain analisis kesintasan. Data klinis diambil dari rekam medis RSUPN Cipto Mangunkusumo sejak Januari 2011-Desember 2016 yang diobservasi selama 2 tahun. Data histopathologik diambil dari Departemen Patologi Anatomi RSUPN Cipto Mangunkusumo yang kemudian dilakukan pulasan imunohistokimia PD-L1.

Hasil: Terdapat hubungan yang bermakna antara ekspresi PD-L1 pada sel tumor dengan 2-year OS ($p=0,023$) dan PFS ($p=0,002$) pada seminoma testis. Tidak terdapat hubungan yang bermakna antara ekspresi PD-L1 pada TILs dengan 2-year OS ($p=0,235$) dan PFS ($p=0,111$). Terdapat hubungan bermakna antara ekspresi PD-L1 pada sel tumor dan ekspresi PD-L1 pada TILs dengan PFS ($p=0,019$). Terdapat hubungan yang bermakna antara stadium dengan 2-year OS ($p=0,010$) dan PFS ($p=0,000$) serta metastasis kelenjar getah bening pada 2-year OS ($p=0,010$) dan PFS ($p=0,000$).

Kesimpulan: Ekspresi PD-L1 pada sel tumor seminoma testis berhubungan dengan OS dan PFS, tetapi hubungan tersebut tidak ditemukan pada TILs.

<hr>

ABSTRACT

Background: The prognosis of testicular seminoma who failed to be cured with conventional therapy is poor. Several recurrent events after chemotherapy were also found. PD-L1 is expressed in various types of malignancy and tumor infiltrating lymphocytes (TILs) and its role is known as a prognostic factor. This study was conducted to determine the role of PD-L1 expression in seminoma in determining overall survival (OS) and progression free survival (PFS).

Materials and Methods: This is a retrospective cohort study with survival analysis. Clinical data were obtained from medical record in RSUPN Cipto Mangunkusumo since January 2011-December 2016 and observed for 2 years. Histopathological data were obtained from Anatomical Pathology Department and PD-L1

immunohistochemistry staining were performed.

Results: A significant correlation between PD-L1 expression in tumor cells with 2-year OS ($p=0,023$) and PFS($p=0,002$) intestinal seminoma was found. No significant correlation between PD-L1 expression in TILs with 2-year OS ($p=0,235$)and PFS ($p=0,111$).We also found significant correlations between PD-L1 expression in tumor cells and TILs with PFS ($p=0,019$). A significant correlation between stage with 2-year OS ($p=0,010$)and PFS ($p=0,000$) and lymph node metastases with 2-year OS ($p=0,010$) and PFS ($p=0,000$)

Conclusion: PD-L1 expression in tumor cells testicular seminoma were significantly correlated with OS and PFS. There were no statistically significant associations between PD-L1 expression in TILs with OS and PFS.