

# Analisis Masalah Terkait Obat pada Pasien Rawat Jalan dengan Hipertensi di RSUD Koja Periode Februari - April Tahun 2019 = Analysis of Drug Related Problem Among Outpatients with Hypertension in Koja District Hospital from February - April 2019

Azizah Nusaibah, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20494112&lokasi=lokal>

---

## Abstrak

### **ABSTRAK**

Pemberian obat untuk pasien hipertensi memerlukan perhatian khusus karena memiliki penyakit penyerta sehingga cenderung mengalami polifarmasi. Penelitian ini bertujuan untuk menganalisis masalah terkait obat dengan domain efektivitas terapi dan reaksi obat yang tidak diinginkan berdasarkan Pharmaceutical Care Network Europe. Metode penelitian ini adalah cross sectional dengan menggunakan rekam medis, resep, dan catatan perawat. Sampel penelitian ini adalah pasien dengan diagnosis primer hipertensi primer dan pasien yang berusia lebih dari 23 tahun. Analisis dilakukan terhadap 114 sampel penelitian. Masalah terkait obat yang paling umum adalah efektivitas terapi dengan persentase 60,00% dan masalah terkait obat lainnya, yaitu reaksi obat yang tidak diinginkan, memiliki persentase 40%. Penyebab masalah tertinggi yang teridentifikasi pada penelitian ini adalah kombinasi obat-obat atau obat-makanan yang tidak tepat termasuk kejadian interaksi obat yaitu 40,00%.

---

### **ABSTRACT**

Giving drugs to hypertensive patients requires special attention because they have comorbidities so they tend to experience polypharmacy. This study aims to analyze drug-related problems with the domain of therapeutic effectiveness and unwanted drug reactions based on Pharmaceutical Care Network Europe. This research method is cross sectional using medical records, prescriptions, and nurse notes. The samples of this study were patients with a primary diagnosis of primary hypertension and patients who were more than 23 years old. The analysis was carried out on 114 research samples. The most common drug-related problems were therapeutic effectiveness with a percentage of 60.00% and other drug-related problems, namely unwanted drug reactions, had a percentage of 40%. The cause of the highest problem identified in this study was an inappropriate drug-drug or drug-food combination including the incidence of drug interactions, which was 40.00%.