

## Risk assesment of air pollution exposure (NO<sub>2</sub>, SO<sub>2</sub>, total suspended particulate, and particulate matter 10 micron) and smoking habits on the lung function of bus drivers in Palembang City

Elvi Sunarsih, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20497479&lokasi=lokal>

---

### Abstrak

#### **ABSTRACT**

Nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), particulate matter (PM<sub>10</sub>), and total suspended particulate (TSP) are the most common and harmful air pollutants to humans. In a short period, air pollution exposure at 5 ppm for 10 minutes in humans causes dyspnea, and when the level is increased to 800 ppm could cause 100% mortality in animals. This study was an analytical study with implementing a cross-sectional design and risk analysis. One hundred subjects were involved in this study. The results showed that the mean value of the non-cancer Hazard Index (HI) for real-time exposure was NO<sub>2</sub>: 1.85; SO<sub>2</sub>: 2.92; TSP: 7.09; and PM<sub>10</sub>: 11.7 (HI value >1). Test for forced vital capacity lung capacity to non-cancer risk estimation of NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> indicated that there was no significant relationship (p-value >0.05). The variable of smoking habit is the most dominant variable (odds ratio [OR] = 12.542) which affects respiratory disorders. The exposure to NO<sub>2</sub>, SO<sub>2</sub>, TSP, and PM<sub>10</sub> in Palembang City bus drivers is considered hazardous to the health of subjects without cancer.