

Pengelolaan Hiu Kejen (*Carcharhinus falciformis* Müller & Henle, 1839) di Perairan Prigi, Jawa Timur = Management of Kejen Sharks (*Carcharhinus falciformis* Muller & Henle, 1839) in Prigi Waters, East Java

Bambang Suseno, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20503859&lokasi=lokal>

Abstrak

Abstrak

Hiu merupakan salah satu kelompok hewan laut yang terancam keberadaan. Hiu yang tertangkap dan didaratkan di PPN Prigi salah satunya adalah hiu kejen (*Carcharhinus falciformis*). Tujuan penelitian adalah 1) menganalisis perikanan hiu kejen yang didaratkan di PPN Prigi; 2) menyusun strategi pengelolaan perikanan hiu kejen yang bertanggung jawab. Penelitian ini menggunakan metode analisis struktur ukuran panjang, hubungan panjang berat, nisbah kelamin, tingkat kematangan gonad, CPUE, MSY, ekologi, ekonomi, kelembagaan dan A'WOT. Hasil tangkapan maksimum lestari di perairan Teluk Prigi sebesar 9.934 kg/tahun. Jumlah hasil tangkapan yang diperbolehkan ditangkap (*total allowable catch*) adalah sebanyak 7.947 kg/tahun. Penangkapan hiu telah membawa dampak secara ekonomis dan ekologis, yang berdasarkan kedua aspek tersebut penangkapan hiu membawa kerugian bagi nelayan. Hasil tangkapan hiu kejen memiliki kisaran panjang total hiu betina 61-162 cm dan jantan 76-189 cm. Pola pertumbuhan tubuh adalah alometrik negatif. Perbandingan jumlah hiu yang tertangkap adalah 0,45:1. Hiu yang didaratkan di PPN Prigi sebagian besar belum mengalami matang gonad (97%). Berdasarkan analisis alternatif strategi pengelolaan yang dapat dilakukan berdasarkan skala prioritas adalah 1) penguatan peran lembaga-lembaga dalam pengawasan pemanfaatan hiu, (2) penetapan pengaturan penangkapan hiu untuk menghindari *overfishing, illegal, unreported,* dan *unregulated fishing*, (3) peningkatan sistem *monitoring, controlling, surveillance* tangkapan hiu.

Abstract

Sharks are a group of marine animals that its existence is threatened. Kejen shark (*Carcharhinus falciformis*) is one of the various types of sharks that are caught and landed on it. The aims of study are 1) analyzing the fisheries of Kejen shark that landed at Prigi fishing port; 2) developing responsible kejen shark resource management strategies. This study uses a long-size analysis method, long weight relationships, sex ratio, gonadal maturity, CPUE, MSY, ecological, economic, institutional and A'WOT. Maximum sustainable catch is insulated by Prigi Bay at 8.619 kg/year and the total allowable catch is 6,895 kg/year. The capture continuously to have an impact on economics and ecology, which are based on those two aspects are carrying losses for fishermen. The catch has a total length range of approximately 61-162 cm for female sharks and 76-189 cm for male sharks. The pattern growth is negative allometric. The ratio of the number of sharks captured is 0.45: 1. The sharks that landed at Prigi fishing port are mostly not from mature gonads (97%). Alternative analysis as a management strategy can be carried out based on priority scales including (1) strengthening the role of institutions in monitoring shark utilization, (2) regulation of shark fishing arrangements to avoid over-fishing, illegal, unreported, and unregulated fishing, (3) improving monitoring systems, controlling, surveillance of shark catches.