

Studi komparasi metode-metode random mutagenesis pada galur aspergillus oryzae dalam meningkatkan produksi asam kojat = Comparative study of random mutagenesis methods in aspergillus oryzae strain in increasing kojic acid production

Tri Hastuti Septiarini, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20504366&lokasi=lokal>

Abstrak

Asam kojat atau kojic acid adalah asam organik yang juga merupakan suatu metabolit sekunder yang dihasilkan oleh berbagai mikroorganisme, salah satunya yaitu Aspergillus oryzae. Dua area utama yang biasanya dikembangkan untuk peningkatan produksi asam kojat adalah pemuliaan galur (strain improvement) dan pengembangan kondisi fermentasi. Hal ini bertujuan untuk mencari metode peningkatan produksi terbaik, yang cepat, murah, efisien, dan tentunya efektif. Random mutagenesis merupakan pendekatan klasik untuk menghasilkan suatu mutan, dimana pada metode ini akan dihasilkan mutasi yang bersifat acak. Sel-sel Aspergillus oryzae yang dipaparkan pada mutagen fisika seperti radiasi sinar UV dan gamma atau pada mutagen kimia seperti N-metil-N-nitro-N-nitrosoguanidin (NTG) dan Etil metan sulfonat (EMS), menunjukkan kemampuannya dalam meningkatkan produksi asam kojat. Beberapa jenis metode mutasi baru seperti Ion Beam Implantation dan Atmospheric and room temperature plasma (ARTP) juga diketahui memiliki efektivitas yang baik dalam upaya meningkatkan produksi produk-produk biologis seperti asam kojat. Komparasi metode random mutagenesis pada galur Aspergillus oryzae dengan penggunaan berbagai jenis mutagen perlu dilakukan untuk memberikan gambaran lebih baik dari tiap metode sehingga dapat membantu peneliti maupun pelajar lainnya dalam mempelajari maupun memilih metode mutagenesis terbaik untuk meningkatkan produksi asam kojat.

<hr>

Kojic acid is an organic acid which is also a secondary metabolite produced by various microorganisms, one of which is Aspergillus oryzae. Two main areas that are usually enhanced for the purpose of increasing kojic acid production are strain improvement and optimization of fermentation conditions. Mutations in kojic acid-producing genes such as the Aspergillus oryzae strain are believed to be effective in increasing the yield of kojic acid. Random mutagenesis is a classic approach for inducing and producing mutants, and in this method, random mutations are produced. Aspergillus oryzae strains that are exposed to physical mutagens such as UV and gamma radiation or to chemical mutagens such as N-methyl-N-nitro-N-nitrosoguanidine (NTG) and Ethyl methane sulfonate (EMS), showed their abilities in increasing the production of kojic acid. Several types of new mutation methods such as Ion Beam Implantation and Atmospheric and room temperature plasma (ARTP) also showed good responses in enhancing the production of biological products such as kojic acid. Comparing these random mutagenesis methods of Aspergillus oryzae with the use of various types of mutagen needs to be done in order to provide a better insight of each method so that it can help researchers and also students in learning and choosing the most suitable method in increasing the kojic acid production.</i>