

## Analisis Pola Pewarisan Bentuk Bunga *Zinnia elegans* yang Tumbuh dari Biji Bunga Tunggal (Single) dan Ganda (Double) = Analysis of Inheritance Pattern on *Zinnia elegans* Flower Shape Growing from Single Layer Flower and Double Layer Flower Seed

Aleta Delviani, author

Deskripsi Lengkap: <https://lib.ui.ac.id/detail?id=20510446&lokasi=lokal>

---

### Abstrak

Penelitian yang bertujuan untuk menganalisis pola pewarisan bentuk bunga *Zinnia elegans* yang tumbuh dari biji bunga tunggal (*single*) dan ganda (*double*) telah dilakukan pada Januari 2020 – Juni 2020. Penelitian dilakukan di 3 lokasi, yaitu Universitas Indonesia, Kelurahan Tanah Baru Depok, dan Kelurahan Grogol Depok. Analisis dilakukan pada 2 generasi tanaman, yaitu pada 237 bunga generasi 1 dan 13 bunga generasi 2. Pengamatan yang dilakukan memerhatikan pola pewarisan jumlah lapisan bunga dan tipe helaian bunga di alam. Hasil yang didapat akan dibandingkan dengan pola pewarisan hukum Mendel, apakah sejalan atau tidak. Berdasarkan hasil penelitian, ditemukan dugaan bahwa lapisan bunga tunggal merupakan tipe yang stabil dan dapat mendominasi tanaman secara alami. Penelitian yang dilakukan juga menduga adanya variasi tipe helaian bunga yang muncul akibat kondisi lingkungan. Tanaman yang berasal dari parental bunga tunggal menunjukkan adanya variasi pola pewarisan, yaitu dapat mengikuti hukum mendel ataupun tidak. Pola pewarisan tanaman yang berasal dari parental bunga ganda belum dapat digambarkan akibat minimnya jumlah tanaman. Pewarisan tipe helaian bunga belum dapat dihubungkan dengan pewarisan Mendel karena informasi tentang tipe helaian bunga sangat terbatas. Pola pewarisan bentuk bunga dari penelitian ini digambarkan dalam bentuk pohon keluarga

(*pedigree*).  
Research to analyze the inheritance pattern on *Zinnia elegans* flower shape was conducted in January 2020 – June 2020. The research was conducted in 3 locations, namely Universitas Indonesia Campus, Kelurahan Tanah Baru Depok, and Kelurahan Grogol Depok. The analysis was conducted on 2 generations of plants, on 237 flowers from first generation and 14 flowers from second generation. The flower shape in nature that has been observed were the number of flower layer and the flower sheet. The results in this research will be compared to Mendelian inheritance pattern, whether follow the rules or not. Based on the results of the study, it was found that a single layer of flower was stable and can dominate the plant naturally. This research also assume that the variation type of flower sheet happen due to environmental conditions. Plants that come from a single layer flower show variation in inheritance patterns, which can either follow the law of Mendel or not. The pattern of the plants derived from double layer flower cannot be describe due to lack of plant quantities. The pattern of flower sheet type cannot be linked to Mendelian inheritance because there is only limited number of studies that provide the information. Inheritance pattern of *Zinnia elegans*'s flower form has been shown on the family tree (*pedigree*).